non-breeding visitor, vagrant

monotypic

Red-breasted Mergansers breed in sub-arctic and n. temperate regions of Eurasia and N America, withdrawing S in winter along the Pacific coasts S to Japan and Baja California (Dement'ev and Gladkov 1952, Cramp and Simmons 1977, AOU 1998) and along Atlantic coasts to Florida and n. Africa. In the Pacific Basin, vagrants have occurred in the Mariana Is (Wiles 2005) and in the Hawaiian Islands in late fall and winter. Half of the records from the Southeastern Hawaiian Islands are from the turn of the 19th century, suggesting that they were formerly much more common than they are at present. All reports have been of females or first-cycle males.

In the *Northwest Hawaiian Islands* two mergansers reported from Sand I., Midway 25 Oct 1981 may have been Red-breasted Mergansers but are unsubstantiated. One on *Laysan* 17-18 Dec 1995 was found dead 21 Jan 1996. A total of 8 substantiated Red-breasted Mergansers have been recorded in the *Southeastern Hawaiian Islands*, on *Kaua'i* (at Hanalei NWR 13 Feb 1995; an adult male reported there in 1998 is unsubstantiated; Engilis et al. 2004), *O'ahu* (one collected in 1901; Bryan 1901a, Henshaw 1902a, whereabouts of specimen unknown; and 1 at Nu'upia Ponds, Mokapu Peninsula Dec 1976, *E* 37:137); *Moloka'i* (Kakahaia NWR 23 Jan 1979; *E* 40:15), *Maui* (Kealia Pond NWR 9 Nov 1985-23 Feb 1986; Engilis et al. 2004), and *Hawai'i I* (two collected near Hilo 8 and 28 Nov 1898 (one of which is USNM 169669; HRBP 5108-5109 of specimen) and one near Pahala in Oct-Nov1902; Henshaw 1900a, 1903). Perkins (1903) mentioned them for "several islands" but we know of records before 1903 only for O'ahu and Hawai'i I.

Acronyms and Abbreviations

Literature cited

Citation: Pyle, R.L., and P. Pyle. 2009. The Birds of the Hawaiian Islands: Occurrence, History, Distribution, and Status. B.P. Bishop Museum, Honolulu, HI, U.S.A. Version 1 (31 December 2009) http://hbs.bishopmuseum.org/birds/rlp-monograph/