

SORA

Porzana carolina

non-breeding visitor, vagrant

monotypic

The Sora breeds from se. Alaska across N America and winters in the s. portion of its breeding range S through n. S America (AOU 1998). It has been recorded as a vagrant to Greenland, Europe (Cramp and Simmons 1980), Clipperton I (Howell et al. 1993), and the *Southeastern Hawaiian Islands*, where there are 3 records. The first record for Hawaii was of a briefly seen but well-described bird in adult winter plumage at the South Pond, Waipi'o Peninsula, *O'ahu*, 10 Feb 1992. Another bird on O'ahu was seen and heard at Ki'i Unit of JCNWR, 20-22 Nov 2003. On *Hawai'i I* an adult was photographed and videotaped at the Cyanotech Lake, Keahole Pt. near Kona 4-22 Sep 2000 (Uyehara 2004; [HRBP](#) 1382, 5713-5717). A report of one from Aimakapa Pond, Hawai'i I, 21 Jan 2009 is unsubstantiated.

[Acronyms and Abbreviations](#)

[Literature cited](#)

Citation: Pyle, R.L., and P. Pyle. 2009. The Birds of the Hawaiian Islands: Occurrence, History, Distribution, and Status. B.P. Bishop Museum, Honolulu, HI, U.S.A. Version 1 (31 December 2009) <http://hbs.bishopmuseum.org/birds/rlp-monograph/>