MARSH SANDPIPER

Tringa stagnatilis

non-breeding visitor, vagrant

monotypic

The Marsh Sandpiper breeds in nc. Asia and winters from Africa through Australia (Dement'ev and Gladkov 1951c, Higgins and Davies 1996). It has reached Europe as a vagrant (Cramp and Simmons 1983), there is a single record for the outer Aleutian Is, Alaska (AOU 1998), and it has been recorded in Micronesia on most island groups E to Kwajalein I in the Marshall Is (Pyle and Engbring 1987, Clapp and Schipper 1990, Wiles 2005, VanderWerf 2006). There are but two records for the Hawaiian Islands: in the *Northwestern Hawaiian Islands*, a first-fall bird occurred on Sand I, *Midway*, 25 Oct-1 Nov 1998 (HRBP 1208-1212), and in the *Southeastern Hawaiian Islands* one was present at the Pouhala Marsh Unit of PHNWR, *O'ahu*, 18 Nov 2002-3 Mar 2003 (HRBP 5554-5559, published *NAB* 57:276).

Acronyms and Abbreviations

Literature cited

Citation: Pyle, R.L., and P. Pyle. 2009. The Birds of the Hawaiian Islands: Occurrence, History, Distribution, and Status. B.P. Bishop Museum, Honolulu, HI, U.S.A. Version 1 (31 December 2009) http://hbs.bishopmuseum.org/birds/rlp-monograph/