

WHITE-RUMPED SANDPIPER

Calidrus fuscicollis

non-breeding visitor, vagrant

monotypic

White-rumped Sandpipers breed in arctic N America and winter in S America (AOU 1998). Vagrants have been recorded in Europe, Russia, Australia and New Zealand, Europe, and California (Dement'ev and Gladkov 1951c, Cramp and Simmons 1983, Higgins and Davies 1996, CBRC 2007), but, outside of the Hawaiian Islands, there are no records known in the central Pacific area. There is one substantiated record for the Southeastern Hawaiian Islands, a bird present at an aquafarm near Kahuku, *O'ahu* 19-28 Sep 2006 ([HRBP](#) 5325-5330 published *NAB* 61:164). A bird observed at Kanaha Pond, Maui 14 Oct 1961 (*E* 22:53) may well have been a White-rumped Sandpiper, as reported, but with few descriptive notes and without photographs we consider it unsubstantiated.

[Acronyms and Abbreviations](#)

[Literature cited](#)

Citation: Pyle, R.L., and P. Pyle. 2009. The Birds of the Hawaiian Islands: Occurrence, History, Distribution, and Status. B.P. Bishop Museum, Honolulu, HI, U.S.A. Version 1 (31 December 2009) <http://hbs.bishopmuseum.org/birds/rlp-monograph/>