RED-THROATED PIPIT

Anthus cervinus

non-breeding visitor, vagrant

monotypic

The Red-throated Pipit breeds across n. Eurasia and Alaska and winters in n. Africa, India, SE Asia, and the Philippine Is (Dement'ev and Gladkov 1954b, Ali and Ripley 1998, Cramp and Simmons 1988, AOU 1998). It is an irregular visitor to California and Baja California in fall, with fewer records for w. North America in spring; in the Pacific it has been recorded on Middleton Is (Alaska), Yap, and Palau (AOU 1998, Wiles 2005, CBRC 2007). In the Hawaiian Islands there are two records, both from *Kure*, *Northwestern Hawaiian Islands*: A first-cycle (PP examination) female was collected 26 Sep 1963 (Clapp and Woodward 1968, Woodward 1972; USNM 493505; HRBP 5462-5464, 5484 of specimen) and a female or first-cycle male was photographed 20-22 Sep 1983 (Pyle 1984; HRBP 0288-0292, 0623).

Acronyms and Abbreviations

Literature cited

Citation: Pyle, R.L., and P. Pyle. 2009. The Birds of the Hawaiian Islands: Occurrence, History, Distribution, and Status. B.P. Bishop Museum, Honolulu, HI, U.S.A. Version 1 (31 December 2009) http://hbs.bishopmuseum.org/birds/rlp-monograph/