## Family PROCRAMPTONOMYIIDAE

The family Procramptonomyiidae was originally proposed based on two species in the genus *Procramptonomyia* Kovalev from the Jurassic deposits of Siberia. Krzemiński (1992) added the genus *Yala* from the Upper Triassic deposits of the Dan River Formation of the eastern United States. Kalugina & Kovalev (1985) claimed that the family is closely related to Cramptonomyiidae (treated in this catalog within the family Pachyneuridae).

Ref.: Kovalev in Kalugina & Kovalev (1985, revision of taxa).

## Genus PROCRAMPTONOMYIA Kovalev

**PROCRAMPTONOMYIA** Kovalev *in* Kalugina & Kovalev, 1985: 119. Type species: *Procramptonomyia sibirica* Kovalev, 1985, by original designation.

incompleta Kovalev *in* Kalugina & Kovalev, 1985: 120. PA: Russia (Siberia) (Middle Jurassic) [C].

sibirica Kovalev in Kalugina & Kovalev, 1985: 119. PA: Russia (Siberia) (Middle Jurassic) [C].

## Genus YALA Krzemiński

**YALA** Krzemiński, 1992 c: 44. Type species: *Yala argentata* Krzemiński, 1992, by original designation.

argentata Krzemiński, 1992 c: 44. NE: USA (Upper Triassic) [C].