

CATALOG OF ENTOMOLOGICAL TYPES IN THE BISHOP MUSEUM.

Orthoptera: Tetrigidae, Trigonopterygidae, Pyrgomorphidae, Acrididae¹

By Gordon M. Nishida²

Abstract. This catalog lists alphabetically under family the following types of Orthoptera deposited at the Bishop Museum: Tetrigidae, 3 holotypes and 2 allotypes; Trigonopterygidae, 1 holotype and 1 allotype; Pyrgomorphidae, 11 holotypes, 2 neotypes and 4 allotypes; and Acrididae, 27 holotypes and 17 allotypes. Associated collection data and information on the condition of the specimens are included.

This catalog completes the listing of orthopteroid types currently deposited in the Bishop Museum collection. The orthopteroid groups previously published were the Tettigoniidae (Nishida 1979b), and Gryllidae (Nishida 1979a).

The approximately 8500 Tetrigidae at the Bishop Museum are housed in 2 Cal Academy insect cabinets. The group has not been comprehensively or recently studied. Material from the Marquesas Is, Micronesia, and Samoa was studied by M. Hebard (1935), C. Willemse (1950), and L. Chopard (1929), respectively, each describing a single species. The 3 species in the type collection are represented by 3 holotypes and 2 allotypes.

The Trigonopterygidae comprise a very small collection of 23 specimens, 21 of which are identified. D.K.McE. Kevan (1966) described the lone species, represented by holotype and allotype in the type collection.

The Pyrgomorphidae were comprehensively studied by D.K.McE. Kevan, who solely described or designated 12 of the 13 species deposited in the type collection. The 12 species are represented by 11 holotypes, 1 neotype and 4 allotypes. Kevan and Y.-K. Chen designated 1 neotype. The Pyrgomorphidae are housed in 1 Cal Academy insect cabinet and consist of approximately 1200 specimens, of which 96% are identified.

The Acrididae type collection consists of 27 holotypes and 17 allotypes, representing 28 species. All were described by F. Willemse with one exception, which was described by D.K.McE. Kevan. The more than 14,000 specimens, housed in 5 Cal Academy cabinets, are 12% identified. Only the subfamilies Oxyinae and Catantopinae have been well studied.

This catalog follows the format outlined by Radovsky et al. (1976).

-
1. This work was partially supported by grant no. BMS 75015971 from the U.S. National Science Foundation to the Bishop Museum.
 2. Department of Entomology, Bishop Museum, P.O. Box 19000-A, Honolulu, Hawaii, 96819, USA.

SPECIES-GROUP ENTRIES

Family TETRIGIDAE

- marquesana** Hebard, **Hydrotetrix** HOLOTYPE ♀ 663
 1935 *Bull. B.P. Bishop Mus.* 114: 124 Fig. 1h-i pin
 MARQUESAS IS: Fatu Hiva I: Omoa Val: Tevaione 518 m 29.VIII.1930 G. Le-Bronnec
 On wet rock by stream
 Filament missing from both antennae. Left hindtarsus missing. ALLOTYPE ♂, 663a, pin, same data as holotype; right antenna missing filament; left hindtarsus missing.
- montana** C. Willems, **Carolinotetrix** HOLOTYPE ♀ 1750
 1950 *Eos, Rev. Esp. Entomol.*, Tomo Extraord.: 348 Fig. 22-24 pin
 MICRONESIA: Caroline Is: Ponape I: Mt Nanalaut 7.III.1936 Z. Ono
 Filament missing from both antennae. Both forelegs missing distad of coxa. Both hindlegs detached, preserved in cellophane.
- samoana** Chopard, **Apterotetrix** HOLOTYPE ♀ 423
 1929 *Insects of Samoa Part 1(2)*: 55 Fig. 50 pin
 SAMOA IS: Tutuila I: Afono Trail 25.IX.1923 O.H. Swezey & G.F. Wilder
 Both antennae missing filament. Right midleg missing. Right hindleg detached, glued to card. Last 2 tarsomeres on right hindleg missing. ALLOTYPE ♂, 423a, pin, SAMOA IS: Savaii I: Safune, 1.V.1924, lowlands to 305 m, E. H. Bryan, Jr.

Family TRIGONOPTERYGIDAE

- curtipennis** Kevan, **Pseudopyrgus** HOLOTYPE ♂ 6740
 1966 *Pac. Insects* 8(2): 392 Fig. 1-3 pin
 MALAYSIA: BORNEO: Sabah: Tenampak 15.II.1959 T.C. Maa
 ALLOTYPE ♀, 6740a, pin & genitalia vial, MALAYSIA: BORNEO: Sabah: Tenampak: 48 km E of Jesselton, 1460 m, 17-18.X.1958, T.C. Maa
 Type labels with locality misspelled as Tenempok.

Family PYRGOMORPHIDAE

- curtipennis** Kevan, **Apodesmoptera** HOLOTYPE ♂ 10716
 1966 *Entomol. Medd.* 34: 399 Fig. 7 Pl. I, fig. b; pl. II, fig. e-g pin &
 PHILIPPINES: Luzon I: Camarines Sur Prov: Mt Isarog 800-1200 m 27.IV.1963 H.M. genitalia
 Torrevillas vial
 Left antenna with 6, right antenna with 9 segments remaining. Left foretarsus missing.
- curvata** Kevan, **Desmopterella** HOLOTYPE ♂ 9184
 1970 *Pac. Insects* 12(3): 616 Fig. 53-55 Pl. XI, fig. A-D pin &
 PNG: NEW GUINEA (NW): Nabire: S Geelvink Bay 0-30 m 2-9.VII.1962 J.L. Gressitt & J. Sedlacek genitalia
 Light trap vial
 Right antenna with 11 segments remaining. Left fore- and midleg missing distad of coxa. Left hindleg missing apical tarsomere. Left wing and tegmen spread. ALLOTYPE ♀, 9184a, pin, same data as holotype; left antenna with 14, right antenna with 10 segments remaining; left wing and tegmen spread.
- elegans** Kevan, **Spinacris** HOLOTYPE ♀ 6632
 1966 *Pac. Insects* 8(1): 5 Fig. 3, 10c, 11c pin &
 PHILIPPINES: Luzon I: Camarines Sur Prov: Mt Isarog 750-850 m 20.IV.1963 H.M. genitalia
 Torrevillas vial
 Both antennae missing distad of scape.
- gracilis** Kevan, **Fijipyrgus** HOLOTYPE ♂ 6809
 1966 *Pac. Insects* 8(2): 399 Fig. 1, 2 pin &

- FIJI: Viti Levu I: Nandarivatu 850 m 8–13.III.1963 C.M. Yoshimoto
Left antenna missing distad of pedicel. Right antenna glued into place at pedicel. Right foreleg missing distad of coxa. Thorax previously separated between prothorax and mesothorax, glued together. Type number published as 6812 in error. genitalia vial
- hystrix** Kevan, **Tarbaleopsis** HOLOTYPE ♂ 6947
1966 *Pac. Insects* 8(3): 725 Fig. 17, 18a–d pin
PNG: NEW GUINEA (SE): S Highlands: Dimifa: SE of Mt Giluwe 2200 m 9.X.1958 J.L. Gressitt
Left antenna with 10, right antenna with 12 segments remaining. Right hindleg missing distal tarsomere. ALLOTYPE ♀, 6947a, pin, same data as holotype.
- microphallica** Kevan, **Stenoxyphula** HOLOTYPE ♂ 10521
1966 *Eos, Rev. Esp. Entomol.* 41(4): 580 Fig. 4, 6 Pl. I, fig. d–g pin & genitalia vial
PNG: NEW GUINEA (NE): Moife: 15 km NW of Okapa 2100 m 7–14.X.1959 T.C. Maa
Both antennae missing distad of scape. Right midleg missing tarsus. Left hindleg missing distal tarsomere. Left tegmen and wing spread. ALLOTYPE ♀, 10521a, pin & genitalia vial, same data as holotype; left midleg missing distad of femur, both hindlegs missing distad of coxa; left tegmen and wing spread.
- minor** Kevan, **Tarbaleopsis** HOLOTYPE ♀ 6945
1966 *Pac. Insects* 8(3): 717 Fig. 11g–h, 13f–g pin
IRIAN: NEW GUINEA (NW): Wissel Lakes: Wagate: Tigi Lake 1700 m 17.VIII.1955 J.L. Gressitt
- novahibernica** Kevan, **Noonacris** HOLOTYPE ♀ 10522
1966 *Pac. Insects* 8(3): 736 Fig. 23g–h, 24c–d (preliminary diagnosis) pin & genitalia vial
1966 *Entomol. Medd.* 34: 392 Fig. 2a–g
PNG: BISMARCK ARCH.: NEW IRELAND: SW: Ridge above "Camp Bishop": 15 km up Kait Riv 250–750 m 13.VII.1956 J.L. Gressitt
Both antennae missing distad of scape.
- oceanicus** Montrouzier, **Truxalis** NEOTYPE ♀ 10628
1969 *Zool. J. Linn. Soc.* 48(2): 191 Pl. 4, fig. A–B pin
PNG: NEW GUINEA (SE): Woodlark I (Murua): Kulumadau Hill 28–30.I.1957 W.W. Brandt
Left hindleg missing distal tarsomere. Right hindleg missing tarsus. Left tegmen and wing spread. Neotype designated by D.K.McE. Kevan and Y.-K. Chen.
- proxima** Kevan, **Tarbaleopsis** HOLOTYPE ♀ 10523
1968 *Proc. R. Entomol. Soc. London* 37(11–12): 157 Fig. 1, 2 Pl. I, fig. 16–17 pin
PNG: NEW GUINEA (NW): Wissel Lakes: Enarotadi 1850–1900 m 30.VII.1962 J. Sedlacek
- schistacra** Kevan, **Meubelia** HOLOTYPE ♂ 7191
1967 *Pac. Insects* 9(3): 443 Fig. 7–10, 25–27, 32, 33, 38 pin & genitalia vial
PHILIPPINES: Mindanao I: Misamis Oriental Prov: Mt Empagatao 1050–1200 m 19–30.IV.1961 H.M. Torre Villas
Left antenna with 14, right antenna with 13 segments remaining. ALLOTYPE ♀, 7191a, pin, same data as holotype, plus rain forest; left antenna with 11, right antenna with 12 segments remaining; right foreleg missing distad of coxa; right midleg missing distal tarsomere.
- stellae** Kevan, **Tarbaleopsis** HOLOTYPE ♂ 6946
1966 *Pac. Insects* 8(3): 719 Fig. 11i–j, 13a–e pin & genitalia vial
IRIAN: NEW GUINEA (NW): Star Mts: Sibil Val 1245 m 18.X–8.XI.1961 S. & L. Quate
Both antennae missing distad of pedicel. Right foreleg missing distal tarsomere. Left midleg missing distal tarsomere. Left hindleg missing tarsal claws.
- sylvaticus** Montrouzier, **Truxalis** NEOTYPE ♀ 9183
1970 *Pac. Insects* 12(3): 601 Fig. 40 Pl. IX, fig. A–B pin
PNG: NEW GUINEA (SE): Woodlark I (Murua): Kulumadau Hill 19–22.III.1957 W.W. Brandt

Left antenna with 12, right antenna with 13 segments remaining. Right midleg, both hindlegs missing distal tarsomere. Left hindleg glued into position. Right tegmen and wing spread. Neotype designated by D.K.McE. Kevan.

Family ACRIDIDAE

- alata** F. Willemse, **Opiptacris** HOLOTYPE ♂ 10673
 1975 *Tijdschr. Entomol.* **118**(6): 139 Pl. 3, fig. 32 pin
 PNG: BISMARCK ARCH.: NEW IRELAND: Manus I: Lorengau 1-75 m 28.VI.1959 J.L. Gressitt
 Pandan
 Both antennae with 13 segments remaining.
- bicingulata** F. Willemse, **Sphaerocranae** HOLOTYPE ♂ 11289
 1974 *Overdr. Public. Natuurhist. Genoot. Limburg* **23**(3-4): 17 Fig. 37-42 Pl. 4, fig. 13-14; pin &
 pl. 5, fig. 17-18 genitalia
 PNG: NEW GUINEA (NE): Feramin 150-120 m 1-6.VI.1959 W.W. Brandt vial
 Date of collection published incorrectly as 7-14.VI.1959. Left antenna missing distad of scape. Left hindleg missing tarsus. Right hindleg missing. Left wing and tegmen spread. Left wing partly missing. ALLOTYPE ♀, 11289a, pin, same data as holotype except 7-14.VI.1959; left antenna missing distad of scape; right antenna with 12 segments remaining.
- bipartita** F. Willemse, **Sphaerocranae** HOLOTYPE ♂ 11288
 1974 *Overdr. Public. Natuurhist. Genoot. Limburg* **23**(3-4): 14 Fig. 27-36 Pl. 3, fig. 11-12; pin
 pl. 4, fig. 15-16
 PNG: NEW GUINEA (NE): Wau 1300 m IV.1968 J. & M. Sedlacek
 Right antenna with 15 segments remaining. ALLOTYPE ♀, 11288a, pin, same data as holotype.
- brunnea** F. Willemse, **Tauchiridea adusta** HOLOTYPE ♂ 11294
 1974 *Overdr. Public. Natuurhist. Genoot. Limburg* **23**(3-4): 35 Pl. 1, fig. 3, 5 pin
 PNG: NEW GUINEA (NE): Kratke Ra: Mt Piora 1650 m 12.VI.1966 O.R. Wilkes
 Grass
 ALLOTYPE ♀, 11294a, pin, same data as holotype; right antenna with 12 segments remaining.
- castanea** Kevan, **Opiptacris** HOLOTYPE ♀ 10357
 1966 *Entomol. Medd.* **34**: 408 Pl. III, fig. a-b pin
 PNG: BISMARCK ARCH.: NEW BRITAIN: Nakanai Mts: Kisiluvu 1050 m 25.VII.1956
 E.J. Ford, Jr.
 Kisiluvu previously published as "Gisiluve." Both antennae missing distad of scape. Right foreleg missing distal tarsomere. Left midleg missing 2 distal tarsomeres. Left hindleg missing. Right hindleg with tarsus detached, glued to label.
- choiseulensis** F. Willemse, **Opiptacris** HOLOTYPE ♂ 10669
 1975 *Tijdschr. Entomol.* **118**(6): 134 Pl. 2, fig. 22-23 pin
 SOLOMON IS: Choiseul I: Kitipi Riv 80 m 17.III.1964 P. Shanahan
 Right antenna missing distad of scape. Right hindleg previously detached, glued into position. ALLOTYPE ♀, 10669a, pin, same data as holotype; left antenna with 15 segments remaining; right antenna missing distad of scape; right foreleg, right hindleg missing distal tarsomere.
- distincta** F. Willemse, **Sphaerocranae** HOLOTYPE ♂ 11287
 1974 *Overdr. Public. Natuurhist. Genoot. Limburg* **23**(3-4): 13 Fig. 21-26 Pl. 2, fig. 6-8; pin &
 pl. 3, fig. 10 genitalia
 PNG: NEW GUINEA (NE): Adelbert Mts 800-1000 m 25.X.1958 J.L. Gressitt vial
 Both antennae missing distad of scape. Right hindleg missing. ALLOTYPE ♀, 11287a, pin, PNG: NEW GUINEA (NE): Adelbert Mts: Wanuma, 25.X.1958, J.L. Gressitt; left antenna missing distad of scape; right antenna with 17 segments remaining; left foreleg missing distal tarsomere; left tegman and wing spread.

- fauroensis** F. Willemse, **Opiptacris bougainvillea** HOLOTYPE ♂ 10672
 1975 *Tijdschr. Entomol.* **118**(6): 138 Pl. 3, fig. 30
 SOLOMON IS: Fauro I 12.IV.1964 P. Shanahan
 Both antennae missing distad of pedicel. All right legs missing distad of coxa. Left midleg missing distal tarsomere. pin
- femorata** F. Willemse, **Opiptacris bougainvillea** HOLOTYPE ♂ 10671
 1975 *Tijdschr. Entomol.* **118**(6): 137 Pl. 3, fig. 28–29
 PNG: SOLOMON IS: Bougainville I: Kokure 690 m 12.VI.1956 E.J. Ford, Jr.
 Right midleg missing distal tarsomere. ALLOTYPE ♀, 10671a, pin, same data as holotype; genitalia
 left antenna with 13, right antenna with 15 segments remaining; right foreleg missing vial
 distal tarsomere.
- genjam** F. Willemse, **Cranae** HOLOTYPE ♂ 11251
 1977 *Tijdschr. Entomol.* **120**(4): 140 Fig. 22–23 Pl. 4, fig. 28
 IRIAN: NEW GUINEA (NW): Genjam: 40 km W of Hollandia 100–200 m 1–
 10.III.1960 T.C. Maa
 Left antenna missing distad of scape, right antenna missing distad of pedicel. Both midlegs
 missing distal tarsomere. vial
- glabra** F. Willemse, **Cranae** HOLOTYPE ♀ 11255
 1977 *Tijdschr. Entomol.* **120**(4): 147 Pl. 1, fig. 6; pl. 2, fig. 15; pl. 3, fig. 23; pl. 5, fig. 33
 IRIAN: NEW GUINEA (NW): Biak I: Kampong Landbouw 50–100 m 27.V.1959 J.L.
 Gressitt pin
- gressitti** F. Willemse, **Pseudocranae** HOLOTYPE ♀ 11285
 1974 *Overdr. Public. Natuurhist. Genoot. Limburg* **23**(3–4): 8 Fig. 9–10 Pl. 1, fig. 2
 PNG: NEW GUINEA (SE): Brown Riv: nr Port Moresby 10 m 5.X.1958 J.L. Gressitt
 Both antennae missing distad of pedicel. Right hind tibia previously detached from femur,
 glued into position. pin
- kevani** F. Willemse, **Cranaella** HOLOTYPE ♂ 11250
 1977 *Tijdschr. Entomol.* **120**(4): 115 Fig. 12–13, 21, 27 Pl. 1, fig. 3
 PHILIPPINES: Mindanao I: Misamis Oriental Prov: Mt Empagatao 1050–1200 m 19–
 30.IV.1961 H. Torrewillas
 Left antenna missing distad of scape. Right antenna with 16 segments remaining. ALLO-
 TYPE ♀, 11250a, pin, same data as holotype; left antenna with 17, right antenna with 16
 segments remaining; right midleg missing distal tarsomere.
- kolombangarae** F. Willemse, **Bumacris pagdeni** HOLOTYPE ♂ 10674
 1975 *Tijdschr. Entomol.* **118**(6): 151 Pl. 4, fig. 37, 43
 SOLOMON IS: New Georgia Group: Kolombangara I: Iriiri 100 m 30.VI.1964 J. &
 M. Sedlacek
 Left tegmen and wing spread. ALLOTYPE ♀, 10674a, pin, same data as holotype; left
 antenna with 13 segments remaining; right antenna missing distad of pedicel; left foreleg
 missing distal tarsomere. pin
- litoralis** F. Willemse, **Pseudocranae** HOLOTYPE ♂ 11286
 1974 *Overdr. Public. Natuurhist. Genoot. Limburg* **23**(3–4): 9 Fig. 11–16 Pl. 1, fig. 3
 PNG: NEW GUINEA (NE): nr Torricelli Mts: Siaute sea level 9–17.XI.1958 W.W.
 Brandt
 Left antenna with 10 segments remaining. Right antenna missing distad of scape. Left
 midleg missing distad of femur. vial
- longipennis** F. Willemse, **Cranae** HOLOTYPE ♂ 11253
 1977 *Tijdschr. Entomol.* **120**(4): 142 Fig. 14–15, 36–37 Pl. 1, fig. 4; pl. 3, fig. 20; pl. 5,
 fig. 31
 PNG: NEW GUINEA (SE): Fly Riv: Olsobip 400–800 m VIII.1969 J. & M. Sedlacek
 Left antenna with 13, right with 14 segments remaining. Left foreleg missing distal tarso-
 mere. Left hindleg missing tarsal claws and arolium. ALLOTYPE ♀, 11253a, pin, same
 data as holotype; left antenna detached distad of pedicel, 14 segments glued back into
 position; right antenna missing distad of pedicel; right foreleg missing distal tarsomere;
 left hindleg missing. vial

- maai** F. Willemse, **Sphaerocranae** HOLOTYPE ♂ 11290
 1974 *Overdr. Public. Natuurhist. Genoot. Limburg* **23**(3-4): 18 Fig. 43-48 Pl. 7, fig. 23-26 pin &
 PNG: NEW GUINEA (NE): Moife: 15 km NW of Okapa 2100 m 7-14.X.1959 T.C. genitalia
 Maa vial
 Right foreleg missing distad of femur. Left tegmen and wing spread. ALLOTYPE ♀,
 11290a, pin, same data as holotype; both antennae with 10 segments remaining; left
 midleg missing tarsus; right midleg missing tarsal claws; right hindleg missing 2 distal
 tarsomeres.
- manokwari** F. Willemse, **Cranae** HOLOTYPE ♂ 11252
 1977 *Tijdschr. Entomol.* **120**(4): 142 Fig. 34-35 Pl. 3, fig. 29 pin &
 IRIAN: NEW GUINEA (NW): Manokwari: Tafelberg Riv 150 m 22.XI.1963 R. Straat- genitalia
 man vial
 Right antenna missing distad of pedicel. Left midleg missing distad of coxa. ALLOTYPE
 ♀, 11252a, pin, same data as holotype; left tegmen spread.
- mundae** F. Willemse, **Bumacris pagdeni** HOLOTYPE ♂ 10675
 1975 *Tijdschr. Entomol.* **118**(6): 151 Fig. 40-41 Pl. 4, fig. 38, 42 pin
 SOLOMON IS: New Georgia Group: New Georgia I: Munda 1-30 m 20.VII.1959 J.L.
 Gressitt
 Right antenna with 16 segments remaining. Left tegmen and wing spread. ALLOTYPE
 ♀, 10675a, pin, same data as holotype; right antenna missing distad of scape; left midleg
 missing distal tarsomere; right tegmen and wing spread.
- nakatae** F. Willemse, **Sphaerocranae** HOLOTYPE ♀ 11293
 1974 *Overdr. Public. Natuurhist. Genoot. Limburg* **23**(3-4): 23 Fig. 57-58 Pl. 2, fig. 9 pin
 PNG: NEW GUINEA (SE): Bisianumu: E of Port Moresby 500 m 8.VI.1955 J.L. Gres-
 sitt
 Primary forest
 Left antenna with 7, right antenna with 9 segments remaining. Left hindleg missing tarsus.
 Left tegmen and wing detached, mounted on card. Left wing partially tattered.
- novageorgica** F. Willemse, **Opiptacris** HOLOTYPE ♂ 10668
 1975 *Tijdschr. Entomol.* **118**(6): 127 Fig. 1-2, 10-13 Pl. 1, fig. 9-10 pin &
 SOLOMON IS: New Georgia Group: New Georgia I: Munda 1-30 m 20.VII.1959 J.L. genitalia
 Gressitt vial
 ALLOTYPE ♀, 10668a, pin, same data as holotype; left antenna with 15, right antenna
 with 14 segments remaining.
- picta** F. Willemse, **Pseudocranae** HOLOTYPE ♂ 11284
 1974 *Overdr. Public. Natuurhist. Genoot. Limburg* **23**(3-4): 7 Fig. 3-8 Pl. 1, fig. 1 pin &
 PNG: NEW GUINEA (SE): Milne Bay 14-23.II.1969 J. Sedlacek genitalia
 Left antenna with 8, right antenna with 16 segments remaining. vial
- poecila** F. Willemse, **Sphaerocranae** HOLOTYPE ♂ 11291
 1974 *Overdr. Public. Natuurhist. Genoot. Limburg* **23**(3-4): 20 Fig. 49-52 Pl. 6, fig. 20-21 pin &
 PNG: NEW GUINEA (NE): Upper Jimmi Val: Tsenga 1200 m 14.VII.1955 J.L. Gres- genitalia
 sitt vial
 Right antenna with 8 segments remaining. Left midleg missing tarsus. Left tegmen and
 wing spread.
- pseudogracilis** F. Willemse, **Sphaerocranae** HOLOTYPE ♂ 11292
 1974 *Overdr. Public. Natuurhist. Genoot. Limburg* **23**(3-4): 21 Fig. 53-56 Pl. 6, fig. 22 pin &
 PNG: NEW GUINEA (NE): Nondugl 2200-2700 m 28.V.1959 C.D. Michener genitalia
 Left antenna with 12 segments remaining. Right antenna missing distad of scape. Left vial
 hindleg detached, glued to determination label. Right hindleg missing distal tarsomere.
- rubra** F. Willemse, **Cranae** HOLOTYPE ♂ 11254
 1977 *Tijdschr. Entomol.* **120**(4): 144 Fig. 38-39 Pl. 5, fig. 32

- PNG: NEW GUINEA (SE): Woodlark I (Murua): Kulumadau Hill 19–22.III.1957 W.W. Brandt pin & genitalia vial
Both antennae missing distad of scape. Both hindlegs missing distal tarsomere. ALLOTYPE ♀, 11254a, pin, same data as holotype; left antenna with 18 segments remaining; right antenna detached at scape, 14 remaining segments glued to determination label.
- tenuis** F. Willemse, **Opiptacris** ALLOTYPE ♀ 10667a pin
1975 *Tijdschr. Entomol.* **118**(6): 125 Pl. 1, fig. 6–7
SOLOMON IS: Guadalcanal I: Gold Ridge 800 m 23.VI.1956 J.L. Gressitt
Left antenna missing distad of pedicel. Holotype deposited British Museum (Natural History), London.
- unicolor** F. Willemse, **Opiptacris** HOLOTYPE ♂ 10670 pin
1975 *Tijdschr. Entomol.* **118**(6): 135 Pl. 2, fig. 24; pl. 3, fig. 25
PNG: SOLOMON IS: Bougainville I: Togerao 600 m 15–21.IV.1968 R. Straatman
Left antenna missing distad of scape. Right antenna with 14 segments remaining. ALLOTYPE ♀, 10670a, pin, same data as holotype except 25–21.IV.1968; right antenna missing distad of pedicel; right foreleg missing tarsus.
- venosa** F. Willemse, **Bumacris** HOLOTYPE ♂ 10676 pin
1975 *Tijdschr. Entomol.* **118**(6): 152 Fig. 42–44 Pl. 4, fig. 46–47
SOLOMON IS: San Cristobal I: Kirakira 30.VII.1960 C.W. O'Brien
Left antenna missing distad of pedicel. Right antenna with 4 segments remaining. Right hindleg missing distal tarsomere. ALLOTYPE ♀, 10676a, pin, same data as holotype except 10.XI.1964, R. Straatman; right antenna with 12 segments remaining.

GENUS GROUP INVENTORY

FAMILY TETRIGIDAE

Apterotettix: samoana
Carolinotettix: montana
Hydrotettix: marquesana

FAMILY TRIGONOPTERYGIDAE

Pseudopyrgus: curtipennis

FAMILY PYRGOMORPHIDAE

Apodesmoptera: curtipennis
Desmopterella: curvata
Fijipyrgus: gracilis
Meubelia: schistacra
Noonacris: novahibernica
Spinacris: elegans
Stenoxyphula: microphallica
Tarbaleopsis: hystrix, minor, proxima, stellae
Truxalis: oceanicus, sylvaticus

FAMILY ACRIDIDAE

Bumacris: pagdeni kolombangarae, pagdeni munda, venosa
Cranae: genjam, glabra, longipennis, manokwari, rubra
Cranaella: kevani
Opiptacris: alata, bougainvillea fauroensis, bougainvillea femorata, castanea, choiseulensis, novageorgica, tenuis, unicolor
Pseudocranae: gressitti, litoralis, picta
Sphaerocranae: bicingulata, bipartita, distincta, maai, nakatae, poecila, pseudogracilis
Tauchiridea: adusta brunnea

LITERATURE CITED

- Chopard, L.** 1929. Orthoptera. *Insects of Samoa*. Part 1(2): 9-58.
- Hebard, M.** 1935. The Dermaptera and Orthoptera of the Marquesas Islands. In: Marquesan Insects—II. *Bull. B. P. Bishop Mus.* **114**: 105-40.
- Kevan, D. K. McE.** 1966. Some Orthoptera-Caelifera from the Philippine, Bismarck and Solomon islands; with a few interesting records from New Guinea and the Moluccas. *Entomol. Medd.* **34**: 375-420.
1966. Further observations on Desmopterini, other than *Desmopterella* Ramme, with descriptions of new species (Orth., Pyrgomorphidae). *Eos, Rev. Espan. Entomol.* **41**(4): 573-96.
1966. Additions to the Verduliini (Orthoptera: Acridoidea: Pyrgomorphidae) from the Philippines. *Pac. Insects* **8**(1): 1-13.
1966. A new genus of Pyrgomorphidae (Orthoptera: Acridoidea) from Fiji. *Pac. Insects* **8**(2): 397-402.
1966. A new genus of Trigonopterygidae (Orthoptera: Acridoidea) from Sabah (North Borneo). *Pac. Insects* **8**(2): 389-96.
1966. The tribe Nereniini: with additions to the Pyrgomorphidae (Orthoptera: Acridoidea) from the South Pacific. *Pac. Insects* **8**(3): 695-758.
1967. New species of *Meubelia* Willemse, 1932, from the Philippines (Orthoptera: Pyrgomorphidae). *Pac. Insects* **9**(3): 437-46.
1968. Some new and little-known species of Pyrgomorphidae (Orthoptera: Acridoidea) from the Pacific and Far East. *Proc. R. Entomol. Soc. London (B)* **37**(11-12): 149-77.
1970. A revision of the Desmopterini (Orthoptera: Acridoidea: Pyrgomorphidae). Part II. *Desmopterella* Ramme, 1941. *Pac. Insects* **12**(3): 543-627.
- Kevan, D. K. McE. & Y.-K. Chen.** 1969. A revised synopsis of the genus *Atractomorpha* Saussure, 1862 (Orthoptera: Pyrgomorphidae) with an account of the African *aberrans*-group. *Zool. J. Linn. Soc.* **48**(2): 141-98.
- Nishida, G. M.** 1979a. Catalog of entomological types in the Bishop Museum. Orthoptera: Gryllidae. *Pac. Insects* **20**(1): 33-36.
- 1979b. Catalog of entomological types in the Bishop Museum. Orthoptera: Tettigoniidae. *Pac. Insects* **20**(1): 27-32.
- Radovsky, F. J., G. A. Samuelson & W. A. Steffan.** 1976. Catalog of entomological types in the Bernice P. Bishop Museum. Introduction. *Pac. Insects* **17**: 1-5.
- Willemse, C.** 1950. On a collection of Orthoptera from the Caroline Islands from the Bernice P. Bishop Museum of Honolulu. *Eos, Rev. Esp. Entomol.*, Tomo Extraord.: 325-61.
- Willemse, F.** 1974. Further notes on Pseudocranae I. Bolivar, 1898, Sphaerocranae F. Willemse, 1972, Salinacris C. Willemse, 1956 and Malua Ramme, 1941 (Orthoptera: Acridoidea: Catantopinae). *Overdr. Public. Natuurhist. Genoot. Limburg* **23**(3-4): 3-32.
1974. Further notes on the genus *Tauchiridea* I. Bolivar, 1918 (Orthoptera: Acridoidea, Oxyinae). *Overdr. Public. Natuurhist. Genoot. Limburg* **23**(3-4): 33-38.
1975. Studies on the acridoid genera *Opiptacris* Walker and *Bumacris* Willemse (Orthoptera: Acridoidea). *Tijdschr. Entomol.* **118**(6): 117-58.
1977. A study on the genus *Cranaella* Ramme (Orthoptera, Acridoidea, Catantopinae). *Tijdschr. Entomol.* **120**(4): 109-20.
1977. A study on the genus *Cranae* Stål (Orthoptera, Acridoidea, Catantopinae). *Tijdschr. Entomol.* **120**(5): 121-52.