

CATALOG OF ENTOMOLOGICAL TYPES IN THE BISHOP MUSEUM Pseudoscorpionida

JoAnn M. Tenorio¹ and William B. Muchmore²

The pseudoscorpion collection in the Bishop Museum, housed in the Arachnology and Ectoparasitology Section of the Department of Entomology, is fairly sizable, considering the secretive nature of these small arachnids and the relatively small number of species recorded in the literature (ca. 2500 species worldwide). Approximately 55% of the collection has been worked, with 135 named species contained in 530 vials (a few on slides); about 37 species are represented by paratypes. The large, unstudied collection numbers some 433 vials; 79% of these represent Australasian (30%), Oriental (24%), and Polynesian (25%) elements.

This catalog includes all holotypes, syntypes and allotypes actually or reportedly deposited in the Bishop Museum. The junior author has examined syntype series, and lectotypes have been designated from most of them. Allotypes are listed as separate entries when the holotype has been deposited elsewhere; when accompanied by a holotype also deposited in Bishop, the allotype holding is simply indicated under the holotype listing. Types that have been transferred to other institutions are listed in a separate section at the end of the main catalog.

Types of 45 species comprising 35 holotypes, 4 lectotypes, 12 allotypes, and 1 syntype are listed as currently deposited in the Bishop Museum; holotypes or lectotypes of 8 species are reported as transferred to other institutions. All except 5 of the species listed herein were described by Max Beier or Joseph C. Chamberlin, both pioneers in the field who made significant contributions to the taxonomy of the world fauna.

The format of this catalog follows Radovsky et al. (1976) and that of previous catalogs. Names are listed alphabetically by families and species or subspecies. Collection information given for type specimens has been corrected or supplemented to the best available knowledge. With the exception of major geographical headings, which may be modified to conform with modern usage and English measurements that are changed to metric equivalents, added information is given in brackets [].

Certain pseudoscorpion species appear to be closely associated with mammals or birds or their nests or with other arthropods. For type specimens listed here that were taken as mammal or bird associates, we have attempted to confirm or correct host identifications for host specimens deposited in Bishop Museum or where other

1. Department of Entomology, Bishop Museum, P.O. Box 19000-A, Honolulu, Hawaii 96819, USA.
2. Department of Biology, University of Rochester, Rochester, New York 14627, USA.

depositories were known. A host identification verified to be correct as published in the original pseudoscorpion description, or one that is corrected or commented on, is marked with an asterisk (*). Those that are not so marked have not been confirmed and are given as originally published. Host confirmations, corrections, or other information relevant to host specimens lodged in Bishop (host numbers prefixed with BM or BBM) were provided by Dr Alan C. Ziegler, Vertebrate Zoologist, Bishop Museum. Confirmation of hosts deposited in the U.S. National Museum was provided by Dr Michael D. Carleton, Assistant Curator, Division of Mammals. For further information on host collections and on problems with field identifications of vertebrate hosts, refer to Tenorio (1976).

Depositories for types and host specimens are abbreviated as follows. American Museum of Natural History, New York—AMNH; British Museum (Natural History), London, England—BMNH; Dominion Museum, Wellington, New Zealand—DMW; Department of Scientific and Industrial Research, Nelson, New Zealand—DSIR; University Museum of Zoology, Cambridge, England—UMZC; U.S. National Museum, Washington, D.C.—USNM.

SPECIES-GROUP ENTRIES

Order PSEUDOSCORPIONIDA

Family ATEMNIDAE

singularis Beier, *Paratemnus*

1965 *Pac. Insects* 7(4): 767-68 Fig. 9

PAPUA NEW GUINEA: New Guinea (SE): Port Moresby 5-6.VI.1955 J.L. Gressitt

Specimen fragmented.

HOLOTYPE ♀ 6264

alcohol

lucifugum Beier, *Cryptocheridium*

1963 *Pac. Insects* 5(1): 51-52 Fig. 1

PENIN. MALAYSIA: Malaya: Batu Caves, Cavern A, floor 29.XII.1959 H.E. McClure

The vial labelled "♂? Typen" contains only a ♀ (with left chela missing). It appears that the ♂, which was also mentioned by Beier in the description, has been lost. Specimens labelled "Paratypen" (2♀, 2TN, 1DN) also in Bishop. Because no single type was designated, all must be considered syntypes. The ♀ type is now designated the lectotype, and the other specimens become paralectotypes.

LECTOTYPE ♀ 3366

alcohol

minutissimum Beier, *Apocheiridium*

1964 *Pac. Insects* 6(2): 312 Fig. 1

PENIN. MALAYSIA: Malaya: Subang 12.XII.1962 [H.E.] McClure

Nest of *Halcyon pileata*, No. M-02066

There is no specimen in the type vial and the holotype is presumably lost. One paratype ♂ (torso) and 1 exuvia in collection.

HOLOTYPE ♀ 3603

alcohol

Family CHELIFERIDAE

exiguus Beier, <i>Papuchelifer</i>	HOLOTYPE ♀	6273
1965 <i>Pac. Insects</i> 7(4): 795–96 Fig. 32		alcohol
IRIAN JAYA: New Guinea (NW): Geelvink Bay, Oransbari [S of Manokwari] 3 m [11.II.1963] [L.P.] Richards		
Ex <i>Alisterus amboinensis</i> *, BBM-NG 22414–15. Host 22414 is in Bishop and is a ♀ <i>A. amboinensis</i> . Host 22415, a ♂, is in AMNH. AMNH identification = <i>Alisterus chloropterus</i> (#799735). No host given originally.		
nigrimanus Beier, <i>Papuchelifer</i>	HOLOTYPE ♂	6272
1965 <i>Pac. Insects</i> 7(4): 793–94 Fig. 30		alcohol
PAPUA NEW GUINEA: New Guinea (SE): Soputa [River, vic. Popondetta] 60 m 1.X.1963 H. Clissold		
Ex kingfisher, prob. <i>Dacelo gaudichaud</i> *, BBM-NG 29769. Field-identified as "same as BBM-29670," a <i>Dacelo gaudichaud</i> ; host spec. not saved by collector.		
orientalis Beier, <i>Hysterochelifer</i>	HOLOTYPE ♀	7422
1967 <i>Pac. Insects</i> 9(2): 358 Fig. 20		alcohol
THAILAND (NW): Chiang Mai Prov, Doi Suthep 1–5.IV.1958 T.C. Maa		
vietnamensis Beier, <i>Tetrachelifer</i>	HOLOTYPE ♀	7423
1967 <i>Pac. Insects</i> 9(2): 364–65 Fig. 24		alcohol
VIETNAM: 13 km W [of] Postal de M'drah 400 m 19.XII.1960 C.M. Yoshimoto		
Nest of <i>Tamiops</i>		
"Yoom" in locality data in original description is a lapsus for 400 m.		

Family CHERNETIDAE

antarcticus Beier, <i>Apatochernes</i>	LECTOTYPE ♂	3873
1964 <i>Pac. Insects Monogr.</i> 7: 116–18 Fig. 1		alcohol
SUBANTARCTIC IS: Campbell I: N.W. Bay 30.XII.1962 K.P. Rennell		
In grass		
Beier placed 1♂ and 1♀ from the above-mentioned locality into a vial and designated them as "Typen"; a long series of others (ca. 130) from the same and other Campbell Island localities and in several vials were called "Paratypen" (1964: 118). It was probably Beier's intention to fix the segregated ♂ and ♀ as holotype and allotype, resp.; however, according to the Code, the entire series must be considered syntypes because no single specimen was designated as the type. We take this opportunity to designate the ♂ of the 2 "Typen" as the lectotype and the ♀ as paralectotype and allotype; all others in the series become paralectotypes. All are in Bishop.		
bougainvillensis Beier, <i>Parachernes</i>	HOLOTYPE ♀	6266
1965 <i>Pac. Insects</i> 7(4): 776 Fig. 16		alcohol
SOLOMON IS: Bougainville I (S): Kokure 690 m 11.VI.1956 [E.J.] Ford, Jr & [J.L.] Gressitt		
Light trap		
Specimen not "trocken" (dry) as stated in description.		
caledonicus Beier, <i>Hebridochernes</i>	HOLOTYPE ♀	3647
1964 <i>Pac. Insects</i> 6(3): 408–09 Fig. 5		alcohol
NEW CALEDONIA: Col d'Amieu 550 m 3.III.1960 J.L. Gressitt		

caledonicus Beier, <i>Nesidiochernes</i>	LECTOTYPE ♂	3646
1964 <i>Pac. Insects</i> 6(3): 406-08 Fig. 3, 4		alcohol
NEW CALEDONIA: Mt. Koghi 28.I.1963 N.L.H. Krauss		
"Typen" 1♂ and 1♀ from the above mentioned locality together in 1 vial; 1♂ and 1♀ designated "Paratypen" are not in Bishop (Beier 1964: 407-08). Although Beier probably intended to fix the ♂ and ♀ "Typen" as holotype and allotype, resp., all the specimens must be considered syntypes according to the Code. We now designate the ♂ of the 2 "Typen" as the lectotype and the ♀ as paralectotype and allotype; the others become paralectotypes.		
carbophilus Beier, <i>Opsochernes</i>		
1966 <i>Pac. Insects</i> 8(2): 370-71 Fig. 6	ALLOTYPE ♀	6736
NEW ZEALAND: South I, Perpendicular Pt 48 km N of Greymouth 10.II.1962 C. Mitchell		alcohol
From Spotted Shag, <i>Stictocarbo punctatus</i> , colony		
Holotype ♂ now deposited in DSIR.		
cornutus Beier, <i>Hebridochernes</i>		
1965 <i>Pac. Insects</i> 7(4): 786-87 Fig. 25	HOLOTYPE ♂	6271
PAPUA NEW GUINEA: New Guinea (NE): Madang 5 m 22.X.1958 [J.L.] Gressitt		alcohol
ALLOTYPE ♀, same data and vial.		
crassus Beier, <i>Megachernes limatus</i>		
1965 <i>Pac. Insects</i> 7(4): 773-74 Fig. 14	HOLOTYPE ♂	6265
IRIAN JAYA: New Guinea (NW): Bokondini, 40 km N [of] Baliem Val 1400 m 19.XI.1961 [L. & S.] Quate		alcohol
Ex unidentified rodents*, BM-NG 337-354. No field identifications available; possibly host specimens not saved by field collectors. Given as <i>Rattus</i> in original description.		
ALLOTYPE ♀, separate vial, Irian Jaya: Star Mts, Sibil Val, 1250 m, 5.XI.1961, L. & S. Quate, ex <i>Melomys</i> cf. <i>platyops</i> *, BM-NG 275.		
gressitti Beier, <i>Hebridochernes</i>		
1964 <i>Pac. Insects</i> 6(3): 409-11 Fig. 6	HOLOTYPE ♀	3648
NEW CALEDONIA: Mt Mou 1200 m 3.II.1963 J.L. Gressitt		alcohol
hawaiensis Simon, <i>Chelifer</i>		
1900 <i>Fauna Hawaiiensis</i> 2: 518 No fig.	1 SYNTYPE ♂	12413
HAWAIIAN IS: Hawaii I: Kona		alcohol
A single ♂ is in the Bishop collection; in addition, at least 4 ♀ of Simon's original series are in the British Museum (Natural History) (see With 1905). By original agreement for distribution of specimens treated in the <i>Fauna Hawaiiensis</i> , the BMNH and UMZC received the 1st two sets and the Bishop Museum the 3rd set, with any remaining going to several other museums; presumably, then, specimens belonging to the type-series are to be found in Cambridge as well as London. Because no type was designated, all are syntypes.		
This species has been designated the type-species of the genus <i>Eumecochernes</i> by Beier (1932).		
It is under this generic name that the specimens are held in the BMNH (personal observation by WBM).		
kanaka Chamberlin, <i>Lamprochernes</i>		
1938 <i>Ann. Mag. Nat. Hist.</i> 11(2): 279 No fig.	HOLOTYPE ♂	2198
MARQUESAS IS: Uapou I: Tekohepu Summit 975 m 28.XI.1931		slide
From dead stipes of <i>Cyathea</i> sp.		
Described in key diagnosis only. Full description and illustrations in Bull. B.P. Bishop Mus., 1939, 142: 210-12, Fig. 2.		
malayanus Beier, <i>Sundochernes</i>		
1963 <i>Pac. Insects</i> 5(3): 510-11 Fig. 2	HOLOTYPE ♂	3465
PENIN MALAYSIA: Malaya: Selangor, Rantau Panjang 8 km N of Klang 28.VI.1961 H.E. McClure & Lim Boo-Liat		alcohol
Nest 191 (B 53365) of <i>Microscelis olivacea</i>		

maoricus Beier, <i>Apatochernes</i>		ALLOTYPE ♀	6739
1966 <i>Pac. Insects</i> 8(2): 375–76 Fig. 10			alcohol
NEW ZEALAND: South I, Banks Penin., Kaituna Val, Scenic Reserve	16.I.1960	C.W.	
O'Brien			
HOLOTYPE ♂ now deposited in DSIR.			
montigenus Beier, <i>Verrucachernes</i>		HOLOTYPE ♀	6267
1965 <i>Pac. Insects</i> 7(4): 779 Fig. 18			alcohol
PAPUA NEW GUINEA: New Guinea (NE): Mt Otto, Kabebe	2200 m	23.VI.1955	[J.L.]
Gressitt			
[Berlesed from] dead tree			
Type #5267 given in original description incorrect. Specimen in poor condition, fragmented.			
monstruosus Beier, <i>Megachernes</i>		LECTOTYPE ♂	6731
1966 <i>Pac. Insects</i> 8(2): 344–46 Fig. 3			alcohol
PHILIPPINES: Palawan I: Mt Mantalingajan	14.IV.1962	M. Thompson	
Nest of <i>Maxomys panglima</i> *. Originally reported as <i>Rattus palawanensis</i> .			
4♂, 2♀, 12N are listed as "Typen" in the original description; only 2♂, 1♀, 11N are in the type vial. We now segregate one of the ♂ and designate it the lectotype, the other specimens becoming paralectotypes.			
navigator Chamberlin, <i>Rhopalochernes</i>		HOLOTYPE ♀	2208
1938 <i>Ann. Mag. Nat. Hist.</i> 11(2): 271–74 Fig. 3			slide
SAMOA: Tutuila taken at quarantine in Hawaii (lot 4827)	4.VIII.1932		
Legs and chelae dissected from body and under separate coverslip on same slide.			
novaeguineae Beier, <i>Nesidiochernes</i>		HOLOTYPE ♀	6269
1965 <i>Pac. Insects</i> 7(4): 780–81 Fig. 20			alcohol
PAPUA NEW GUINEA: New Guinea (NE): Maprik	160 m	15.X.1957	[J.L.] Gressitt
novaeguineae Beier, <i>Sundochernes</i>		HOLOTYPE ♀	6268
1965 <i>Pac. Insects</i> 7(4): 779–80 Fig. 19			alcohol
PAPUA NEW GUINEA: New Guinea (SE): Mt Giluwe	2550 m	28.V.1963	[J.] Sedlacek
papuanus Beier, <i>Hebridochernes</i>		HOLOTYPE ♂	6270
1965 <i>Pac. Insects</i> 7(4): 784–86 Fig. 24			alcohol
PAPUA NEW GUINEA: New Guinea (SE): Daradae, nr Javarere, Musgrove Riv	100 m	5.X.1958	[J.L.] Gressitt
ALLOTYPE ♀, same data and vial			
Original report gives elevation as 100 mm.			
philippinus Beier, <i>Megachernes</i>		HOLOTYPE ♀	6730
1966 <i>Pac. Insects</i> 8(2): 343–44 Fig. 2			alcohol
PHILIPPINES: Mindanao I: Dapitan Pk	XI.1959	L.W. Quate	
Ex prob. <i>Rattus</i> sp., BBM-PI 571 Field identified as <i>Rattus</i> sp.; host spec. prob. saved by field collector but its present whereabouts unknown			
Type with 2TN and 2DN in same vial.			
tropicus Beier, <i>Eremochernes</i>		HOLOTYPE ♀	7421
1967 <i>Pac. Insects</i> 9(2): 357–58 Fig. 19			alcohol
THAILAND (NW): Chiang Mai Prov, Doi Suthep	1–5.IV.1958	T.C. Maa	
vietnamensis Beier, <i>Megachernes</i>		HOLOTYPE ♂	7420
1967 <i>Pac. Insects</i> 9(2): 355–56 Fig. 16			alcohol
VIETNAM: [Thac Da Tan La]	1550 m	14.IX.1960	C. Yoshimoto & B. Feinstein
Ex <i>Niviventer bukit</i> * (USNM #321392–94) and <i>Rattus sikkimensis</i> * (USNM #321228–29).			
Originally reported as <i>Rattus niviventer</i> and <i>R. s. sladeni</i> .			
ALLOTYPE ♀, same data and vial. "Dalat" given as locality in orig. description.			
yapensis Beier, <i>Smeringochernes (Smeringochernes)</i>		ALLOTYPE ♀	2554
1957 <i>Insects Micronesia</i> 3(1): 42–43 Fig. 24, 25c			alcohol
MICRONESIA: Yap: Ruul Distr	[VII–]VIII.1950	[R.J.] Goss	
HOLOTYPE ♂ in USNM.			

zonatus Beier, <i>Systellochernes</i>	ALLOTYPE ♀	3874
1964 <i>Pac. Insects Monogr.</i> 7: 119-20 Fig. 2		alcohol
SUBANTARCTIC IS: Campbell I: St. Col-Lyall Saddle 3-5.XII.1961 [J.L.] Gressitt		
Mixed moss		

HOLOTYPE ♂ in DMW.

Family CHTHONIIDAE

howarthi Muchmore, <i>Tyrannochthonius</i>	HOLOTYPE ♀	11335
1979 <i>Pac. Insects</i> 20(2-3): 187-190 Fig. 1-5		slide
HAWAIIAN IS: Hawaii I: Hawaii Volcanoes Natl. Park, Petroglyph Cave 700 m 13.VII.1976 F.G. Howarth		
Dark zone of cave		
obscurus Beier, <i>Austrochthonius</i>	ALLOTYPE ♀	6733
1966 <i>Pac. Insects</i> 8(2): 364-65 Fig. 2		alcohol
NEW ZEALAND: South I: Banks Penin, Kaituna Val, Scenic Reserve 16.I.1960 C.W. O'Brien		
Two specimens, "Typen" ♂ and ♀, described. Because neither was designated the holotype, both are syntypes, according to the Code. We now designate the ♂ (presently deposited in DSIR) as the lectotype and the ♀ in Bishop as paralectotype and allotype.		
philippina Beier, <i>Morikawia</i>	HOLOTYPE ♀	6728
1966 <i>Pac. Insects</i> 8(2): 340-42 Fig. 1		alcohol
PHILIPPINES: Mindanao I: Bukidnon, Mt Katanglad 1480 m 27-31.X.1959 L. Quate & C. Yoshimoto		
<i>Morikawia</i> Chamberlin has been shown to be synonymous with <i>Paraliochthonius</i> Beier (Muchmore 1972). It is not yet clear whether this species belongs in <i>Paraliochthonius</i> or some other genus.		
webbi Muchmore, <i>Sathrochthonius</i>	HOLOTYPE ♂	12592
1982 <i>Pac. Insects</i> 24(2): 156 Fig. 1, 2		slide
AUSTRALIA: Queensland: Holy Jump Lava Cave, 25 km E [of] Warwick 2.III.1980 J.A. Webb		
On bat guano		
zonata Beier, <i>Morikawia</i>	HOLOTYPE ♂	3645
1964 <i>Pac. Insects</i> 6(3): 403-05 Fig. 1		alcohol
NEW CALEDONIA: Mt Koghi 26-30.I.1963 N.L.H. Krauss & C.M. Yoshimoto Berlese		
See remarks under <i>philippina</i> Beier, <i>Morikawia</i> regarding generic placement.		

Family GARYPIDAE

albus Beier, <i>Geogarypus</i>	HOLOTYPE ♂	3464
1963 <i>Pac. Insects</i> 5(3): 508-10 Fig. 1		alcohol
PENIN. MALAYSIA: Malaya: Selangor, Rantau Panjang 8 km N of Klang 1960 H.E. McClure & Lim Boo-Liat		
Nest (#65) of <i>Microscelis olivacea</i>		
ALLOTYPE ♀, same data except "nest 62."		
australianus Muchmore, <i>Anagarypus</i>	HOLOTYPE ♂	12593
1982 <i>Pac. Insects</i> 24(2): 160 Fig. 1-6		slide
AUSTRALIA: Great Barrier Reef: Nymph I 23.IX.1967 H. Heatwole		
Under stone		

heatwolei Muchmore, <i>Anagarypus</i>	HOLOTYPE ♀	12594
1982 <i>Pac. Insects</i> 24(2): 162 Fig. 7	slide	
AUSTRALIA: Western Australia: Barrow I 11.II.1977 H. Heatwole		
Under stone on a bare rocky headland		
marquesianus Chamberlin, <i>Geogarypus (Geogarypus)</i>	HOLOTYPE ♀	2197
1939 <i>Bull. B.P. Bishop Mus.</i> 142: 208–10 Fig. 1	slide	
MARQUESAS IS: Uahuka I: Putatauuua, Vaipee Val 268 m 21.IX.1929 A.M. Adamson		
From dead banana leaves		
sagittatus Beier, <i>Geogarypus (Indogarypus)</i>	HOLOTYPE ♀	6262
1965 <i>Pac. Insects</i> 7(4): 764–65 Fig. 7	alcohol	
IRIAN JAYA: New Guinea (NW): Ifar, Cyclops Mts 300 m 20.VI.1959 [J.L.] Gressitt		

Family HYIDAE

vietnamensis Beier, <i>Stenohya</i>	HOLOTYPE TN	7419
1967 <i>Pac. Insects</i> 9(2): 345 Fig. 4	alcohol	

VIETNAM: Dalat, 6 km S [of] 1400–1500 m 9.VI–7.VII.1961 N.R. Spencer

Family OLPIIDAE

cyclopium Beier, <i>Xenolpium</i>	HOLOTYPE ♀	6261
1965 <i>Pac. Insects</i> 7(4): 763–64 Fig. 6	alcohol	

IRIAN JAYA: New Guinea (NW): Cyclops Mts, nr Kota Nica 420 m 28.XII.1961 L. & S. Quate

In cave with pteropodid bat*, BM-NG 490. Field-identified as "Dobsonia"; apparently host specimen(s) not saved by collectors.

Family PSEUDOCHIRIDIIDAE

triquetrum Beier, <i>Pseudochiridium</i>	HOLOTYPE ♂	6263
1965 <i>Pac. Insects</i> 7(4): 765–67 Fig. 8	alcohol	

IRIAN JAYA: New Guinea (NW): S side of L. Sentani, Ajappo 31.XII.1961 L.W. Quate

Nest of prob. apodid bird, BBM-NG 586-620. Field identified as "swift"; host specimens apparently not saved by collector.

ALLOTYPE ♀, same data and vial.

Family SYARINIDAE

philippinense Beier, <i>Alocobisium</i>	HOLOTYPE ♀	6729
1966 <i>Pac. Insects</i> 8(2): 342 No fig.	alcohol	

PHILIPPINES: Palawan I: Tarumpitao Pt 21.V.1958 H.E. Milliron

In jungle duff

Specimen in poor condition, fragmented.

Family TRIDENCHTHONIIDAE

loricata Beier, <i>Ditha</i>	HOLOTYPE ♂	6260
1965 <i>Pac. Insects</i> 7(4): 757–58 Fig. 2	alcohol	

IRIAN JAYA: New Guinea (NW): Vogelkop, Oransbari 12.XII.1962 H. Holtmann

Under bark

ALLOTYPE ♀, same data and vial.

HOLOTYPE TRANSFERRED TO OTHER INSTITUTIONS

Family CHTHONIIDAE

obscurus, Austrochthonius: see general listing

zealandica Beier, *Tyrannochthoniella*

1966 *Pac. Insects* 8(2): 367-68 Fig. 4

HOLOTYPE ♀ now in DSIR.

zealandicus Beier, *Austrochthonius*

1966 *Pac. Insects* 8(2): 363-64 Fig. 1

HOLOTYPE ♀ now in DSIR.

Family GARYPIDAE

lineatus Beier, *Synsphyronus (Maorigarypus)*

1966 *Pac. Insects* 8(2): 368-69 Fig. 5

HOLOTYPE ♂ now in DSIR.

Family CHERNETIDAE

carbophilus, Opsochernes: see general listing

maoricus, Apatochernes: see general listing

obrieni Beier, *Apatochernes*

1966 *Pac. Insects* 8(2): 374-75 Fig. 9

HOLOTYPE ♀ now in DSIR.

zealandicus Beier, *Nesidiochernes*

1966 *Pac. Insects* 8(2): 372-73 Fig. 8

HOLOTYPE ♀ now in DSIR.

FAMILY AND GENUS-GROUP INVENTORY

Alocobisium: philippinense

Anagarypus: australianus, heatwolei

Apatochernes: antarcticus, maoricus, obrieni

Apocheiridium: minutissimum

Atemniidae

Austrochthonius: obscurus, zealandicus

Cheiridiidae

Chelifer: hawaiiensis

Cheliferidae

Chernetidae

Chthoniidae

Cryptocheiridium: lucifugum

Ditha: loricata

Eremochernes: tropicus

Garypidae

Geogarypus: albus, marquesianus, (Indogarypus) sagittatus

Hebridochernes: caledonicus, cornutus, gressitti, papuanus

Hyidae

Hysterochelifer: orientalis

Indogarypus: subg., see Geogarypus

Lamprochernes: kanaka

Maorigarypus: subg., see Synsphyronus

Megachernes: limatus crassus, monstrosus, philippinus, vietnamensis

Morikawia: philippina, zonata

Nesidiochernes: caledonicus, novaeguineae, zealandicus
 Olpiidae
 Opsochernes: carbophilus
 Papuchelifer: exiguus, nigrimanus
 Parachernes: bougainvillensis
 Paratemnus: singularis
 Pseudochiridiidae
 Pseudochiridium: triquetrum
 Rhopalochernes: navigator
 Sathrochthonius: webbi
 Smerringochernes: yapensis
 Stenohya: vietnamensis
 Sundochernes: malayanus, novaeguineae
 Syarinidae
 Synsphyronus (Maorigarypus): lineatus
 Systellochernes: zonatus
 Tetrachelifer: vietnamensis
 Tridenchthoniidae
 Tyrannochthoniella: zealandica
 Tyrannochthonius: howarthi
 Verrucachernes: montigenus
 Xenolpium: cyclopium

LITERATURE CITED

- Beier, M.** 1932. Pseudoscorpionidea II. Subord. C. Cheliferinea. *Tierreich* **58**: 1-294.
 1957. Pseudoscorpionida. *Insects Micronesia* **3**(1): 1-64.
 1963. Pseudoscorpione von den Batu-Höhlen in Malaya. *Pac. Insects* **5**(1): 51-52.
 1963. Pseudoscorpione aus Vogelnestern von Malaya. *Pac. Insects* **5**(3): 507-11.
 1964. Insects of Campbell Island. Pseudoscorpionidea. *Pac. Insects Monogr.* **7**: 116-20.
 1964. Some further nidicolous Chelomathi (Pseudoscorpionida) from Malaya. *Pac. Insects* **6**(2): 312-13.
 1964. Pseudoscorpione von Neu-Caledonien. *Pac. Insects* **6**(3): 403-11.
 1964. Further records of Pseudoscorpionidea from the Solomon Islands. *Pac. Insects* **6**(4): 592-98.
 1965. Die Pseudoscorpioniden Neu-Guineas und der benachbarten Inseln. *Pac. Insects* **7**(4): 749-96.
 1966. Über Pseudoscorpione von den Philippinen. *Pac. Insects* **8**(2): 340-48.
 1966. Zur Kenntnis der Pseudoscorpioniden-Fauna Neu-Seelands. *Pac. Insects* **8**(2): 363-79.
 1967. Pseudoscorpione vom kontinentalen Südost-Asien. *Pac. Insects* **9**(2): 341-69.
Chamberlin, Joseph C. 1938. New and little-known false-scorpions from the Pacific and elsewhere (Arachnida—Chelonethida). *Ann. Mag. Nat. Hist. Ser. 11, 2*: 259-85.
 1939. New and little-known false scorpions from the Marquesas Islands (Arachnida: Chelonethida). In: Marquesan Insects III. *Bull. B.P. Bishop Museum* **142**: 207-15.
Muchmore, William B. 1972. The pseudoscorpion genus *Paraliochthonius* (Arachnida, Pseudoscorpionida, Chthoniidae). *Entomol. News* **83**: 248-56.
 1979. The cavernicolous fauna of Hawaiian lava tubes. 11. A troglobitic pseudoscorpion (Pseudoscorpionida: Chthoniidae). *Pac. Insects* **20**(2-3): 187-90.
 1982. A new cavernicolous *Sathrochthonius* from Australia (Pseudoscorpionida: Chthoniidae). *Pac. Insects* **24**(2): 156-58.
 1982. The genus *Anagarypus* (Pseudoscorpionida: Garypidae). *Pac. Insects* **24**(2): 159-63.
Radovsky, F.J., G.A. Samuelson & W.A. Steffan. 1976. Catalog of entomological types in the Bernice P. Bishop Museum. Introduction. *Pac. Insects* **17**: 1-5.
Simon, Eugène. 1900. Arachnida. p. 443-519. In: Sharp, David, ed., *Fauna Hawaiensis*. Vol. 2.
Tenorio, JoAnn M. 1976. Catalog of entomological types in the Bernice P. Bishop Museum. Subclass Acari. *Pac. Insects* **17**(1): 7-46.
With, C.J. 1905. On Chelonethi, chiefly from the Australian region, in the collection of the British Museum, with observations on the "coxal sac" and on some cases of abnormal segmentation. *Ann. Mag. Nat. Hist. Ser. 7, 15*: 94-143.