

DIPTERA: PSYCHODIDAE OF SOUTH GEORGIA¹By D. A. Duckhouse²

Abstract: One species of Psychodidae—*Psychoda parthenogenetica* Tonnoir is reported from South Georgia.

***Psychoda parthenogenetica* Tonnoir**

Psychoda severini parthenogenetica Ton., 1940, *Trans. Soc. Brit. Ent.* **7**: 53–54.

Psychoda parthenogenetica: Duckhouse, 1962, *Trans. R. Ent. Soc. Lond.* **114**: 428.

SOUTH GEORGIA: 1 ♀, Grytviken Pen., King Edward Cove, 12.XII.1963, on grass, along ditch, sea level, H. B. Clagg; 5 ♀♀, Fortuna Bay, 8.I.1964, from tussock grass, sea level to 150 m, Clagg.

This widely distributed but certainly not cosmopolitan, parthenogenetic species, has been recorded from Europe, Japan, Korea, New Zealand, southern Australia, and from a number of other remote islands besides South Georgia, including Kerguelen and Macquarie (Womersley 1937), Campbell (Quate 1964) and the Juan Fernandez group (Satchell 1952).

LITERATURE CITED

Quate, L. W. 1964. Insects of Campbell Island. Diptera: Psychodidae. *Pacif. Ins. Monogr.* **7**: 280–88.

Satchell, G. H. 1952. Los insectos de las Islas Juan Fernandez 8. Psychodidae (Diptera). *Rev. Chile. Ent.* **2**: 105–7.

Womersley, H., 1937. Diptera. Rep. *B. A. N. Z. Antarc. Res. Exped.* 1929–31, **4**: 59–79.

¹Results of fieldwork supported by grants (G-23720, GA-166) to Bishop Museum from the Office of Antarctic Programs, U. S. National Science Foundation.

²University of Adelaide, South Australia.