

NEW PALMS FROM FIJI

By

MAX BURRET

BERNICE P. BISHOP MUSEUM

OCCASIONAL PAPERS

VOLUME XI, NUMBER 4

HONOLULU, HAWAII

PUBLISHED BY THE MUSEUM

May 25, 1935

NEW PALMS FROM FIJI

By MAX BURRET

Most of the plants described in this paper were collected by Dr. A. C. Smith in 1933 and 1934, with the aid of a Bishop Museum Fellowship in Yale University. The first set of specimens is deposited in the herbarium of Bernice P. Bishop Museum.

For convenience, one species from Samoa is described here.

Genus PRITCHARDIA Seemann and H. Wendland

Pritchardia thurstoni F. von Mueller and Drude.

Fiji: Fulanga, A. C. Smith 1230, on limestone cliff in lagoon. Local names are *masai* and *viu*. The species is very common on the small islets in the lagoon; it has previously been collected only once in a wild state.

Genus EXORRHIZA Beccari

Exorrhiza smithii, species nova.

Palma 4-8 m alta; caudex robustus curvatus annulatus; frondis vagina cylindrica, ad apicem extus fusco-leproso-maculata; petiolus evolutus, inter minores, visum fragmentum inferius 20 cm longum, ad basin 4 cm, superne 2.5 cm in diam. metiens, marginibus acutis, supra concavus, subtus rotundato-triangularis, in modo jam descripto fusco-maculatus; segmenta numerosa, regulariter disposita, directione haud variantia, a rhachi sat patentia sed superne cernua, sat robusta, linearia, haud curvata, maxima visa 62 cm longa, 2.4 cm lata, basi leviter reduplicata, superne sensim attenuata et acuminata, longe anguste producta, in apices 2 perangustos profunde fissa, 3-nervia, costa media et nervo secundario fere aequivalido dextra sinistra percursa, nervis tertiaris numerosis densis conspicuis, subtus iisdem nervis et tertiaris densis, prominentibus, squamulis minutis fuscis dense praeditis, costa media inferne paleis badiis linearibus in dimidia parte affixis notata; segmenta superiora sensim latitudine et longitudine decrescentia, sensim longe anguste acuminata; spadix infra frondes, scopiformis, duplo-(forsan triplo?) ramosus; rami 2 visi ramulos floriferos 5 gerentes, applanati, breves; ramuli rhachi brevi inserti, basi bractea late ovata breviter subito acuminata suffulti, pro rata tenues, 20 cm circ. longi, i. s. longitudinaliter dense striati, glabri, apice in filum flexuosum sat longum producti; florum glomeruli in spira densa dispositi, in alveolis sat profundis; axis margine triangulariter nonnihil producto, superne curvato, acuto; glomeruli ad maximam ramorum partem 3-flori, bracteis conspicuis late ovatis, flore ♀ intermedio, ♂ binis superpositis, ad ramorum apicem floribus ♂ tantum; flores ♂ non visi, omnes delapsi, ♀ parvi, ovati; sepala ovata; petala ovata, apice breviter triangulari valvata; fructus ignoti.

Fiji: Vanua Levu, Thakaundrove, eastern buttress of Mount Ndi-keva, altitude 900 meters, June 5, 1934, A. C. Smith 1879, type.

It is a species related to *E. wendlandiana* Beccari, but with the spadix much smaller, the branchlets much shorter and thinner, and the leaf segments straight rather than falcate. The segments of the new species are perhaps not so deeply split at the apex, and on the under surface bear more densely congested brown scales. I have not seen the male flowers of *E. smithii*, which are probably smaller than those of *E. wendlandiana*. The following new Samoan species is also of this alliance.

***Exorrhiza vaupelii*, species nova.**

Frondis rhachis superne triangularis, in marginibus crispato-floccoso-fusco-tomentosa, in lateribus et subtus atro-leproso-punctulata; segmenta numerosa, regulariter disposita, directione haud variantia, majora visa 80 cm circ. longa, usque 3.4 cm lata, textura firma, basi modice, sed conspicue reduplicata, linearia, haud curvata, maxima latitudine in parte inferiore sita, dein sensim attenuata et longe anguste sensim acuminata, in apices 2 angustos aequales fissa, 3-nervia, costa media valde prominente, ad basin modo descripto fusco-tomentosa, subtus inferne paleis badiis linearibus numerosis densissimis notata, nervis lateralibus praecipue ad basin supra plicato-prominentibus, minus superne; nervi tertiarii numerosi, utrinque conspicui, subtus squamulis atris densis praediti; segmenta superiora longitudine et latitudine decrescentia.

Samoa: Savaii, south of Mangaloa, altitude 1200 meters, F. Vaupel 605, type, in Herb. Berlin.

E. vaupelii resembles *E. smithii* in its linear leaflets which are not at all curved, with very numerous minute dark scales on the lower surface. I have little doubt as to the proper genus; the only other genus to be compared is *Chinostigma*. *C. oncorhynchum* Beccari has not the dense minute scales on the under surface of the leaflets. I have not seen the leaflets of *C. samoense* H. Wendland, but the above-mentioned minute scales are apparently lacking, and the rachis is said to be quite glabrous. *C. powellianum*, from the description, is very close to *C. samoense*.

Genus **VEITCHIA** H. Wendland

***Weitchia joannis* H. Wendland.**

Fiji: Vanua Levu, Thakaundrove, southern slope of Korotini Range, below Navitho Pass, 300-650 meters, A. C. Smith 575. In Seemann's "Flora Vitiensis" this palm is mentioned as being very common throughout Fiji. In the wet forest of Vanua Levu, however, it is rare, having been observed only twice by the collector. It appears to be the tallest palm of the region, the present specimen being over 100 feet in height.

Genus **BALAKA** Beccari**Balaka longirostris** Beccari.

Fiji, without exact locality, D. Jeoward, in 1894 (non vidi).

This palm has been described by Beccari only from fruits, without perianth. It has the largest fruit as yet known in the genus *Balaka*. The plant below described is possibly identical with *B. longirostris*, but from the description and figure of the fruit of that species, I believe the new species to be quite distinct. The fruits of *B. longirostris* are 32 mm long; the fruits of our new species are 40 mm and more long, having more excavate sides, the rostrum much broader, and the sclerosoms of the surface quite distinct, in which characters it differs from the description and figure of Beccari's species. The new species may be known as:

Balaka macrocarpa, species nova.

Palma 4-8 m alta, caudice gracili, 4-6 cm in diam. metiente; frondes ex fragmentis inter maiores, circ. 2 m longae; vagina ut petiolus et rhachis fusco-furfuracea, demum minute fusco-leprosa, vagina cylindrica, robusta, dorso i. s. 6 mm in diam. metiens, fibris densis numerosis percursa, apice sat breviter contracta; petiolus 13 cm circ. longus, robustus, basi 1.2 cm, apice 1 cm fere latus, supra canaliculatus, marginibus acutis, dorso late angulato-rotundatus; segmenta prob. sat numerosa, infima ambo angusta, linearia, 8-15 mm lata, rhacheos prob. mediae 2 visa in margine superiore 38-41 cm longa, leviter sigmoidea, alterum nervis primariis 2 percursum, superne 10.5 cm latum, inferne sensim angustatum, ad basin 3.2 cm in diam., rhachin secus 6 cm latum, apice oblique lobulato-praemorsum, alterum costa media unica, ad apicem 7.2 cm latum, basi nonnihil angustiore, rhacheos fragmentum apicale summis segmentis ambobus latissimis, rhacheos 21.5-24.5 cm amplectentibus, apice ipso oblique praemorso, obtuse triangulariter lobulato, 17 cm lato, nervis primariis 8 percursis, segmento inferne sequente viso leviter cuneato, 35 cm longo, superne 4.5 cm lato, nervo primario uno percurso; spadix 3 dm circ. longus; rhacheos fragmentum ut rami ramulique i. s. sat acute longitudinaliter angulosi, densissime conspicue granulosi; rami primarii 2 visi, basi bractea brevi late rotundata suffulti, ramulis 2-3; ramuli fructiferi paulo curvati, ad fructuum insertiones modice flexuosi, inter robustiores, in dimidia circ. parte 2-2.5 mm in internodiis in diam. metientes, superne sensim angustati, visi 15-20 cm longi, fructibus spiraliter dispositis; ex insertionibus florum glomeruli ad magnam ramuli partem 3-flori, flore ♀ intermedio, ♂ binis dextra sinistra insertis, superne 2-flori, flore femineo et masculo juxtapositis, demum flore ♂ solitario; fructus basi bracteis erectis late rotundatis calyculum humilem, 1 mm altitudine paulo superantem formantibus cincti, magni, i. v. aurantiaci, i. s. brunneo-flavidi, 4.2 cm longi, quadrangulares, 1.5 cm in diam. metientes, angulis longitudinalibus acutis valde prominentibus, lateribus inter angulos nonnihil excavatis, pericarpii rugis perpaucis, sed plerumque prominentibus transverse vel plus minus oblique pertensis, rostrati; rostrum excentrice productum, crassum, angulariter conicum, 13 mm longum; perianthium cupuliforme, laxe accumbens;

calyx delapsus; corolla 12 mm alta, petalis late rotundatis, apice breviter triangulari haud producto, obsoletō, longitudinaliter dense nervoso-striatus; epicarpium sclerosomatibus densis lineolatis conspicuis; pericarpium carnosum, siccum pro rata sat evolutum; endocarpium pergamentaceum; semen acute 4-angulare, 22 mm longum, 6-7 mm in diam. metiens, rostrum conspicue anguste acuminatum, 8-9 mm longum; raphe ab apice in angulo longe decurrentis, ramis in quoque latere 4, dorso sat numerosis descendenteribus, parce furcatis vel anastomosantibus; embryo basilaris; albumen aequabile; flores ♂ haud perfecte evoluti visi 5 mm longi, ovato-oblongi, fusco-leproso-maculati; calyx 2.5 mm altus, sepalis basi umbonatis, superne nervoso-striatis, apice late rotundatis, ciliatis; petala oblonga costata; stamina 8; filaments filiformia; antherae lineares, basi sagittatae, apice incisae, dorso ad basin affixa; pistillodium inferne dilatatum, superne in stylum productum, longitudine staminum.

Fiji: Vanua Levu, Thakaundrove, Natewa Peninsula, Uluingala, altitude 600-820 meters, June 15, 1934, A. C. Smith 2007, type.

This slender and attractive palm, known as *niuniu*, was seen only once. Several trees were growing close together in the shade of large forest trees, on a sharp well-drained slope.

Balaka cuneata, species nova.

Palma 2-5 m alta; caudex 1.5 cm circ. in diam., annulis, ut videtur, sat dissitis; frondis vagina fusco-furfuracea, in latere petiolo opposito producta; petiolus circ. 20 cm longus, sordide flavidoto-tomentosus atque fusco-lepidoto-leprosus; rhachis 65 cm fere longa, furfuracea et fusco-leprosa, segmentis utrinsecus 9; segmenta sat regulariter disposita, infimum cujusque lateris quam reliqua aliquanto minus, praecipue angustius, media circ. 23 cm longa, 7 cm lata, in margine superiore producta, margine superiore oblique lobulato-prae-morso, apicalia confluentia, nervis primariis 3 percursa, rhacheos 4.3-5.5 cm amplectentia, in margine superiore 18 cm longa, 6-7 cm in diam., oblique lobulato-prae-morsa, reliqua nervo primario unico percursa, omnia supra nitentia, glabra, subtus fere opaca, ad margines et in segmentis apicalibus ambobus ad nervos secundarios fusco-furfuracea, papryacea, cuneiformia, ad basin haud curvata, vix reduplicata, costa basin versus haud curvata; spadix 35 cm fere longus, duplicato-ramosus; pedunculus 13 cm longus, applanatus, decidue fusco-furfuraceus, spathe inferioris annulo ad basin, superioris 3 cm altius sito atque bractea semiannulari 4 cm supra eum notatus; rhachis vix flexuosa, ramis primariis ramificatis 4 atque simplicibus reliquis 7, infimis 2 ramulos fructiferis 4 (secundo) vel plures (infimo), sequentibus 3-2 gerentibus; ramuli fructiferi majores usque 16 cm longi visi, plurimi breviores, omnes superne sensim attenuati, vix curvati, haud flexuosi; fructus ex glomerulis in spira perlaxe dispositis i. s. cum periantho 1.3 cm longi, usque 6 mm lati, oblique nonnihil rostrati, inferne 4-angulares, sclerosomatibus densissime gibbosí, nigri; perianthium sat explanatum, tenué; sepalum late rotundatum; petala lata, margine ciliata, longitudinaliter dense nervoso-striata, apice triangulare valvata.

Fiji: Vanua Levu, Thakaundrove, hills between Vatukawa and Wainingio Rivers, Ndrekeniwai Valley, altitude 200-500 meters, November 24, 1933, A. C. Smith 577, type; Natewa Peninsula, hills south of Natewa, altitude 400-600 meters, A. C. Smith 1950.

The species, locally known as *mbalaka*, is near *B. seemannii*, (H. Wendland) Beccari, from which it differs by the cuneate leaflets, which, with the costa, are not curved at the base. The spadix is much more ramified than that of *B. seemannii*. The other species of the genus known from Fiji have much shorter and broader deltoid leaflets. *B. longirostris* Beccari has fruits twice as long and thick.

Genus VITIPHOENIX Beccari

Subgenus 1. EUVITIPHOENIX, subgenus novum

Frondis segmenta apice praemorsa; bracteae florem ♀ et fructum cingentes explanatae angustissimae obsoletae.

This subgenus includes the following species: *V. filifera* (H. Wendland) Beccari, *V. vitiensis* (H. Wendland) Burret, *V. picker-ningii* (H. Wendland) Burret, *V. pauciflora* (H. Wendland) Burret, and :

Vitiphoenix (Euvitiphoenix) smithii, species nova.

Palma 16 m alta; frondes 3 m longae; vaginae apex visus, extus ut petiolus et rhachis praeципue subtus cinereo- (an in vivo fusco)-tomentosus et minute dense vel sat dense fuso-lepidotus; petioli fragmentum a vagine apice 16.5 cm longum visum subtus rotundato-triangulare, marginibus acutis, supra marginem versus paulo excavatum, dimidium secus leviter convexum, basi 18 mm latum, ad apicem vix angustius; segmenta numerosa, regulariter disposita, maxima visa 41.5 cm longa, usque 5 cm lata, leviter sigmoidea, basin versus conspicue, superne leviter curvata, maxima latitudine ad dimidię vel modice infra sita, inferne nonnihil fere contracto-angustata, ad basin angusta, 8 mm circ. lata, paulo sed conspicue reduplicata, superne nonnihil angustata, ad apicem oblique lobulato-praemorsum 1.8 cm tantum lata, supra nitentia, subtus opaca, sub lente saepe minutissime gibbosa, costa media unica, subtus basin versus paleis badiis linearibus densis praedita, nervis marginalibus validis, inferne subtus fuso-lepidoto-punctulatis, nervis minus validis sat numerosis, ulterioribus nullis; segmenta superiora sensim longitudine et latitudine decrescentia, fere rectilinearia, maxima latitudine in dimidia circ. sita, inferne nonnihil angustata, reduplicata sed etiam apicem versus conspicue angustata, apice oblique lobulato-praemorsa, apicalia ambo quam sequentia conspicue latiora, 2 cm circ. lata, 10 cm in margine superiore longa, apicem versus paulo attenuata, apice transverse lobulato-praemorsa; spadix infra frondes, 1 m longus, 4-plo multiramosus; rhachis et rami visi glabri; rhachis et rami primarii applanato-subtriangulares, rami (visi) basi bractea haud producta, anguste lineolata; ramus primarius maximus visus 30 cm longus; rami ramulique i. s. longitudinaliter lineolati, ramuli ultimae ordinis plurimi, breves et perbreves, pauciflori, longissimus visus 6.5 cm longus, majores 2.5 cm minores numerosi 1 cm vix aequantes, fructibus 2-1, omnes inter tenuiores, in internodiis 1 mm circ. in diam, metientes, superne attenuati, ad nodos leviter flexuosi; florum glomeruli inferiores 3-flori, ♀ medio, ♂ binis juxtapositis, ad apicem ramulorum 2 prob. masculi; fructum insertiones

obliqui, quam axis duplo latiores, bracteis explanatis; fructus i. s. cum perianthio 12 mm longus, ambitu clavato-oblongus, superne usque 7 mm in diam., apice haud producto, breviter conico, extus glabri, longitudinaliter dense fibroso-lineolati, sclerosomatibus nullis; perianthium fructiferum 6-7 mm altum, cupuliforme, ore 5-6 mm in diam. metiente; calyx scutelliformis, sepalis late rotundatis, nervoso-striatis; corolla 3-plo fere altior, petala nervoso-striata, late rotundata, imbricata, apice breviter trianguli valvata; mesocarpium fibris numerosis, endocarpium tenuis; semen ambitu oblongum, leviter claviforme, teres, 11 mm longum, superne 5.5 mm in diam.; raphae linearis, totum latus sequens, ramis ex rhapheos parte superiore et dorso paralleliter descendantibus, parce anastomosantibus; embryo basilaris; albumen aequabile; staminodia in perianthii basi perpaucia, visa 2-3 sat late dentiformia; flores ♂ nondum perfecte evoluti visi 4 mm longi, ambitu ovato-oblongi; calyx 2 mm altus, sepalis late rotundatis ciliatis; petala sat crassa; stamina numerosa, filamentum filiforme, demum prob. sat evolutum; antherae oblongo-lineares, basi sagittatae, apice incisae, dorso ad basin affixae; pistillodium inferne fusiforme, superne filiforme, longitudine staminum.

Fiji: Kandavu, hills above Namalata and Ngaloa Bays, altitude 200-400 meters, October 16, 1933, A. C. Smith 162, type.

V. smithii differs from the other known Fijian species, except *V. vitiensis*, which is described only from leaves, by having the spadix more branched, with the thin branchlets much shorter and with few fruits. The leaflets of the new species are conspicuously attenuate towards the apex, and the rachis is tomentose below in a very characteristic peculiar manner, whereas in *V. vitiensis* the superior leaflets do not taper towards the end, the rachis being only leprose rather than tomentose.

Subgenus 2. ACMOPHOENIX, subgenus novum

Frondis segmenta apice acuta; bracteae florem ♀ et fructum cingentes surgentes, conspicuae, calyculum formantes.

Although the following two species differ considerably from those of *Euvitiphoenix*, differences in the fruits, seeds, and flowers do not seem sufficient to necessitate a new genus.

Vitiphoenix (Acmophoenix) petiolata, species nova.

Palma 15-35 m alta; vagina ut petiolus et rhachis subtus furfuracea atque punctato-fusco-furfuracea; petiolus in fragmendo ab apice vaginæ 25 cm circ. longo visus, supra excavatus, marginibus acutis, subtus rotundato-triangularis; segmenta infima jam sat dense sese sequentia, omnia regulariter disposita, prob. dimidiae circ. rhacheos lanceolata, 55 cm longa, 4 cm lata, maxima latitudine in tertia inferiore sita, inferne sensim angustata, basi sat angusta, leviter sed conspicue curvata, superne sensim attenuata et acuminate, inaequaliter bifida, apice superiore multo longiore, anguste, sat longe acuminato, inferiore brevi, textura rigida, costa media subtus basin versus paleis anguste linearibus, i. s.

tortis dense praedita, nervis marginalibus sat robustis, minus validis dextra sinistra pluribus, paulo conspicuis, tenuibus obscuris, segmenta superiora longitudine et latitudine decrescentia, linear-lanceolata, superne attenuata, apice breviter bifida; spadix 3-plo ramosus, pedunculatus, rhachis, ut rami et ramuli inferne longitudinaliter angulosi, i. s. dense, ramuli superne minus dense gibbos; rami primarii ramosi sat numerosi; ramuli floriferi 10-12 cm longi, in internodiis dimidiae partis i. s. 1.5 mm in diam., raro paulo crassiores, superne leviter flexuosi, usque ad apicem fere glomerulis in spira laxa dispositis, 3-floris, ♀ internodio, ad apicem tantum 2-floris, masculis ad summum apicem breviter fere filiformes; flores ♂ 7 mm circ. longi, oblongi; calyx i. s. patens, sat explanatus, 4 mm in diam. metiens, sepalis late rotundatis; petala oblonga; stamina numerosa, circ. 50; filamenta filiformia, quam antherae breviora; antherae lineares, 3 mm longitudine superantes, basi sagittatae, apice incisae, dorso basi affixae; pistillodium filiforme, apice bifidum, an semper?, longitudine fere statim; bracteae florem ♀ cingentes surgentes, conspicuae; flos ♀: sepala late rotundata, petala late rotundata, apice triangularis; fructus inter minores angustiores, cum perianthio i. s. 1.8 cm longus, 0.6 cm in diam. metiens, ambitu oblongo-lanceolatus, breviter sed conspicue rostratus, extus sclero-somatibus densissime sat longe lineolatus; pericarpium i. s. 0.5 mm in diam. vix superans; mesocarpium fibris numerosis percursum; endocarpium tenue; semen ambitu oblongo-lanceolatum, 1 cm fere longum, 4 mm in diam., teres, apice basique rotundatum; albumen aequabile; embryo basilaris; perianthium cupulare, 5 mm altum, 5 mm in diam.; calyx humilis, sepalis late rotundatis, extus nervosis; petala late rotundata, apice breviter triangulari sed tamen quam margines conspicue altiore; staminodia pauca, plus minus late dentiformia.

Fiji: Vanua Levu, Mbua, southern slope of Mount Seatura, altitude 500 meters, April 28, 1934, A. C. Smith 1687, type.

This palm, growing in dense wet forest, is one of those known as *niuniu*. Compared to *V. sessilifolia* (below described), the petiole in *V. petiolata* is quite conspicuous, the fruit is much smaller, narrower, and rostrate at the top, the calyx of the male flowers is patent and explanate, the triangular valvate top of the petals of the fruiting perianth is conspicuously longer than the margins, whereas the top of the petals in *V. sessilifolia* is shorter than the protracted margins.

Vitiphoenix (Acmophoenix) sessilifolia, species nova.

Palma 10 m alta; frondis vagina cylindrica, ut rhachis praecipue subtus punctulato-fusco-leprosa atque plus minus decidue furfuracea; frondes quam homo adultus modice longiores, superne cernuae, segmentis regulariter dispositis, majoribus, ut videtur levissime sigmoideis; petiolus nullus vel fere, basi 1.8 cm latus, ut rhachis inferior supra excavatus, dorso rotundato-triangularis, circ. 7 cm longus, sed segmento secundo utrinsecus 10-11 cm ab infimo remoto; segmenta infima angusta, dimidiae prob. rhacheos lanceolata, 50 cm circ. longa, 5 cm lata, rigida, inferne nonnihil sat subito angustata, superne longe acuminata, acuta, an bifida apicibus plus minus destructis incerte, utrinque viridia, costa media paleis subtus carens, et marginalibus nonnihil prominentibus, minus validis dextra sinistra paucis, paulo conspicuis, tenuibus nullis, frondis fragmenti apicalis segmenta longitudine et latitudine decrescentia, anguste linear-lanceo-

lata, sensim acuminata, acuta; spadix 3(4?)-plo ramosus, pedunculatus, ramis primariis inferioribus longe pedunculatis pluribus; rami primo fusco-furfuracei, longitudinaliter plus minus angulosi; ramuli 9, 5-6 cm longi, in dimidio in internodiis 1 mm diam. paulo superantes, superne sensim attenuati, ad apicem sat tenues, superne ad fructuum insertiones levissime flexuosi; florum glomeruli usque ad ramulorum apicem 3-flori, flore ♀ intermedio, ♂ juxtapositis, ♀ bracteis surgentibus conspicuis circumdato; flores ♂ 6 mm longi, ambitu elliptico-oblongi, apice rotundati; calyx corollae arcte accumbens, 2.5 mm altus, 3 mm in diam. metiens, sepalis late rotundatis, minute ciliatis, inferne crassis, umbonatis; petala oblonga, crassa, apice obtusa; stamna numerosa, circ. 40; filamenta filamentosa, quam antherae breviores; antherae lineares, 2-3 mm fere longae, basi sagittatae, apice plerumque incisae, dorso ad basin affixae; pistillodium ad basin ipsam leviter fusiforme, superne filiforme, longitudine staminum, ut videtur rarissime haud evolutum; flores ♀ juveniles: sepala late rotundata, minute ciliata, inferne crassa, umbonata; petala rotundata, apice triangulari valvata; ovarium ambitu ovato-oblongum, stigmatibus 3 triangularibus; staminodia ut videtur plus minus in annulum connata; fructus maturus aurantiacus, i. s. cum perianthio 2.2 cm longus, ambitu oblongus, 1 cm in diam., apice rotundatus, apice mammilli, perbreviter latissime conico imposito, epicarpio minute longitudinali secus lineolato; pericarpium 1 mm in diam. metiens, mesocarpio fibris numerosis totam longitudinali secus percurso, endocarpio tenuiter pergamaceo, fragili; semen ambitu oblongum, 11 mm longum, 6 mm in diam. metiens, apice basique rotundatum, rhabpe totum latus secus extensa linearis, ramis ex rhabpe parte superiore pluribus atque dorso descendentibus parce furcatis et anastomosantibus; embryo basilaris; albumen aequabile; perianthium fructiferum cupuliforme, 7 mm altum; calyx sat explanatus, humilis, 7 mm in max. diam. metiens, sepalis late rotundatis, longitudinaliter nervoso-striatis; petala latissime rotundata, apice valvato breviter triangulari, quam margines rotundati breviore; staminodia perpaucia, latissime dentiformia.

Fiji: Vanua Levu, Thakaundrove, Yanawai River region, Mount Kasi, altitude 300-430 meters, May 10, 1934, A. C. Smith 1784, type.

Like the preceding, this palm grows in dense forest and is known as *niuniu*.

Genus GONIOSPERMA, genus novum

Spadix patenter ramosus; florum glomeruli spiraliter dispositi, ad majorem ramorum partem 3-flori; femineo intermedio, ♂ binis dextra sinistra juxtapositis, apicem ramorum versus ♂ geminis, glomeruli superficialiter fere axi inserti, sed axi infra in forma bracteae triangulariter conspicue producta; flores ♂ symmetrici; calyx sepalis late rotundatis, nonnihil imbricatis; corolla petalis valvatis, circ. oblongis, apice obtusis; stamna 6 introrsa; filamenta applanata, inter longiora, apice geniculatim nonnihil inflexa; antherae lineares, thecis intus dense parallelis, connectivo dorsili angusto, dorso modice infra dimidiata insertae, basi incisae, apice paulo emarginatae vel

incisae; pistillodium cylindricum sat robustum, quam stamina brevius; flores ♀ : sepala nonnihil imbricata rotundata; petala imbricata, apicibus triangularibus valvata; staminodia prob. 6, dentiformia; gynoecium oblongum, stigmatibus 3 triangularibus; ovulum totam longitudinem secus affixum; fructus inter maiores, globosi, apice modice excentrico mamillato, carnosus; epicarpium nitens, fere laeve, minute haud dense gibbosum; mesocarpium carnosum, fibris nullis; endocarpium tenuiter osseum, prob. longitudinaliter 4-angulare; semen longitudinem secus acute 4-angulare, apice oblique excavato-obtusum; rapha in angulo totum latus secus pertensa, linearis; embryo basilaris; albumen homogeneum. Palma altitudinis mediorum, foliis petiolatis, regulariter pinnatisectis; segmenta nervo primario unico, apice acuta.

The type of the genus is designated as *G. vitiense*. The genus is related to *Physokentia* Beccari, based on *P. tete* Beccari, of which I have not seen material. From description, the flowers of that species seem very similar to those of the new genus. The pinnae are premorse at the apex rather than acute. According to the description and figure, the seed of *Physokentia* is without distinct acute longitudinal edges, typical of the new genus. The rapha of *Physokentia* is not described.

The new species here described is certainly very near *G. thurstonii*, possibly even identical with it. The earlier species is known only from the fruit, so that a comparison of it with *G. vitiense* is hardly possible. In my opinion it is preferable to describe a new species rather than to establish a genus on a possible mixture of two species.

Goniosperma vitiense, species nova.

Palma caudice 4 m alto, petiolo 30 cm longo, lamina 3 m longa, segmentis utrinsecus circ. 40; rhachis prob. circ. dimidiae frondis humilius triangularis, lateribus nonnihil excavatis, subtus levissime rotundata, utrinque ut videtur fusco-furfuracea atque dense fusco-leproso-unctulata; segmenta linearis-lanceolata, fere rectilinearis, apice vix falcata, visa regulariter disposita, opposita vel fere, sequentia 6.5 cm circ. dissita, a rhachi patentia, sed leviter surgentia, 75 cm circ. longa, 6 cm circ. lata, maxima latitudine ut videtur in dimidia parte circ. sita, inferne sensim sed conspicue angustata, basi conspicue reduplicata, superne sensim attenuata, apice acuto haud producto, textura sat rigida, supra glabra, subtus praecipue inferne in nervis validioribus paleis badiis plus minus oblongis praedita et in nervis nervulisque minutissime fusco-lepidota, supra in nervis validioribus fere plicata; spadix infra frondes 50 cm circ. longus; rhachis applanato-subtriangularis, decidue breviter furfuracea; rami ramulique basi bractea modice triangulariter producta suffulti; rami primarii

6 mm latum, ut rami ramulique trichomatibus paleaceis, lacerato-ciliatis, i. s. pallide fuscis vel fere cinereis dense furfuraceum; rhachis et rami ramulique longitudinaliter angulosi; rhachis 26 cm longa, haud flexuosa; rami ramulique basi bractea conspicua plus minus triangulariter ovata suffulti, basi in angulo interiore nonnihil callosi, rami primarii inferiores 8 ramosi, 7 reliqui simplices, infimi 2 ramulos 7 emittentes, rami imi pedunculus 8.5 cm longus, rhachis 7 cm; ramuli vix curvati, haud flexuosi, longiores 18 cm, breviores circ. 12 cm longi, inter tenuiores, floriferi in parte validiore, inferiore circ. 2 mm, in dimidia 1 mm (in internodiis) in diam. metientes, superne angustati, apice tenui acuto; florum glomeruli nonnihil foveolati, spiraliter inferne sat laxe, superne densiuscule dispositi; fovea margine inferiore nonnihil, late rotundato-producta; glomeruli in ramulorum parte inferiore dimidiata vix aequante

2-flori, flore ♀ et ♂ juxtapositis, superne ♂ tantum; flores ♂ parvi, symmetrici, fere ovati, 2.5 mm longi, apice rotundati; calyx 1.5 mm fere altus, sepalis i. s. dorso carinatis, margine late rotundato ciliato, extus nervoso-striatis; petala valvata, sat firma, apice obtusa, dorso longitudinaliter conspicue nervoso-striata; stamina 6; filamenta libera, filiformia, sat evoluta, apice geniculato-inflexa; antherae oblongae, 1 mm longae, introrsae, in medio dorso affixae, basi incisae, apice integrae; pistillodium longitudine staminum, robustum, cylindricum, apice nonnihil capitatum; flores ♀ bracteis surgentibus, foveae labrium vix superantibus; sepala rotundata, late imbricata, longitudinaliter ut petala nervoso-striata; petala imbricata, apice triangulariter valvata; pistillodium obovato-oblongum, stigmatibus 3 triangularibus; ovulum parietale.

Fiji: Taveuni, October 2, 1881, C. Weber 112, type in herb.
Berlin.

The species is said to be widely distributed in the hills of Taveuni,
generally below 300 meters.