

Notes on the Naturalized Flora of the Hawaiian Islands.

By CHARLES N. FORBES.

THE following plants are mainly species not recorded in Hillebrand's Flora,¹ the majority of them having probably become naturalized since 1886. The only record of the naturalized plants which we have had since that time is an addition of a few species in an article by Heller.² In order to make these notes complete these species are again mentioned in this paper. It is the intention to continue the notes at irregular intervals as material accumulates.

The naturalized flora occupies the lower valleys and plains, the number of indigenous plants in these localities at present being insignificant. This flora is rapidly encroaching on the native forest, especially where its conditions are changed by fires or grazing. Introduced weeds appear along new trails through the native forest in from two to three weeks in places where it would be impossible to find them before. In the unforested sections of the islands, as a large proportion of the slopes of Haleakala on Maui, naturalized plants occur to the very summit (10,000 feet). A few of the naturalized plants as *Crepis japonica* are an exception to the rule, in that they occur in the dense forest and are rarely found on the plains; and perhaps a few of the plants considered as indigenous may belong in this category.

¹ W Hillebrand. *Flora of the Hawaiian Islands*, 1888.

² A. A. Heller. *Minnesota Botanical Studies, Bulletin No. 9*, pp. 760-922. A portion of Heller's collections are in the Herbarium of the Bishop Museum.

FILICALES.

Ceropteris calomelanos* (L.) Und.Acrostichum calomelanos* L., Sp. Pl., ii, 1753, 1072.*Gymnogramma calomelanos* Kaulf., Enum., 1824, 76.*Ceropteris calomelanos* (L.) Und., Torr. Cl., 1902, 29, 632.

The silver fern is well established on all the islands, generally occurring along roadsides and irrigating ditches.

Ceropteris calomelanos* var *chrysophylla* Link.Gymnogramme chrysophyllum* Kl., Enum., 1824, 74.*Ceropteris chrysophylla* Link, Sp., 1841, 143.

The gold fern has the same distribution as the former but is less common. On Hawaii it was observed growing in the steam cracks at Kilauea.

HYDROCHARITACEAE.

Halophila ovata* Gaud.Halophila ovata* Gaud., Freyc. Voy. Bot., 1826, 430, t. 40. f.*Halophila ovalis* Hook., Fl. Tas., 1860, ii, 45.

Miss M. Reed showed me a specimen of this plant which she said was identified by Dr. W. A. Setchell. It occurs in brackish ponds in the vicinity of Honolulu.

LEMNACEAE.

Lemna minor* L.Lemna minor* L., Sp. Pl., 1753, 970.

The duck weed is common in ponds and taro patches on all the islands.

CHENOPodiACEAE.

Amaranthus spinosus* L.Amaranthus spinosus* L., Sp. Pl., 1753, 991.

A rather common weed on Oahu, probably occurring on other islands as well.

C. N. F., no. 1060. O. Nuuanu Valley.

POLYGONACEAE.

Rumex acetocella* L.Rumex acetocella* L., Sp. Pl., 1753, 338.

The sorrel occurs on all the islands and at all elevations. First reported from Kauai by Heller.

Rumex patentia L.*Rumex patentia* L., Sp. Pl., 1753, 333.

A few specimens of this species were observed at Nahiku Landing, E. Maui. The plant also occurs as a weed in cane fields on Hawaii.

C. N. F., no. 265. M.

PORTULACACEAE,

Portulaca lutea Sol.*Portulaca lutea* Sol., Ex Forst. f. Pl. Escul., 1786, 72. Seem. Fl. Vit., 9.*Portulaca flava* DC., Prod., 1843, iii, 355.

This species occurs as a rather rare shore plant, probably on all the islands. This museum has specimens collected on Marcus, Midway and Laysan islands by W. A. Bryan, while it has been reported from Necker island by W. K. Fisher³ and from Hawaii by Dr. Guppy.⁴ It also occurs on Oahu near Kaena Point, and should be considered as indigenous.

C. N. F., no. 1650. O.

CARYOPHYLLACEAE.

Drymaria cordata (L.) Willd.*Holosteum cordatum* L., Sp. Pl., 1753, 88.*Drymaria cordata* Willd., Roem & Schultes, Syst. Veg., 1819, 5, 406.

Observed by Heller on Kauai. Occurs on all the islands.

Polycarpon tetraphyllum L.*Polycarpon tetraphyllum* L., Sp. Pl., 1753, 881.

Observed at Makawao, E. Maui.

C. N. F., Aug. 1910. M.

Tissa marina (L.) Britton.*Arenaria rubra var marina* L., Sp. Pl., 1753, 423.*Spergularia salina* Presl., Fl. Cech., 1819, 95.*Buda marina* Dumort, Fl. Belg., 1827, 110.*Spergularia media* A. Gray, Man., 1867, 5, 95.*Tissa marina* Britton, Bul. Torr. Club, 1889, 16, 126.*Buda marina var minor* S. Wats., A. Gray, Man., 1890, Ed. vi, 90.

Near the beach in various places between Diamond Head and Koko Head on Oahu.

C. N. F., no. 1501. O.

³W. K. Fisher. Bul. U. S. Fish Commission, vol. xxiii, pt. iii, 1903, p. 807.

⁴H. B. Guppy. Observations of a Naturalist in the Pacific, vol. ii, p. 555.

RANUNCULACEAE.

Ranunculus parviflorus L.*Ranunculus parviflorus* L., Sp. Pl., 1763, Ed. ii, 780.

In the forest below Kaalapuuwale on the slopes of Hualalai on the island of Hawaii, occurs a species of *Ranunculus* which I refer to the above. This diffuse plant with minute yellow flowers and hispid carpels covers many acres; mainly in the dead Koa forest, although some specimens were found in virgin forest. This place is also a locality for *R. mauiensis*, the two species often growing close together; while within a mile can be found *R. hawaiiensis*, the other native species. My determination is based on the description found in Britton and Brown's Flora of the Northern States and Canada.

C. N. F., no. 275. H.

ANONACEAE.

Anona cherimolia Mill.*Anona cherimolia* Mill, Gard. Dic. Ed. viii, 1768, n. 5.

The cherimolia is well established on Hawaii, especially in the drier localities.

C. N. F., no. 365. H.

CRUCIFERAE.

Brassica campestris L.*Brassica campestris* L., Sp. Pl., 1753, 666.

Rather common on Oahu, also occurring on the other islands.
C. N. F., March 18, 1909. Moanalua Valley, Oahu.

Raphanus sativus L.*Raphanus sativus* L., Sp. Pl., 1753, 669.

Rare, to be expected on all the islands. Mentioned by Hillebrand as occurring on Kauai.

C. N. F., April 6, 1909. Moanalua Valley, Oahu.

Sisymbrium officinale (L.) Scop.*Erysimum officinale* L., Sp. Pl., 1753, 660.*Sisymbrium officinale* Scop., Pl. Carn. Ed. ii, 1772, 2, 26.

The hedge mustard was first observed at Puuwaawaa on Hawaii, and later at various places on that island.

C. N. F., no. 9. H.

ROSACEAE.

Rosa sp.

A small flowered multi-petaled rose is well established on Hawaii and in certain places on Maui. On Hawaii it is known as the volcano rose, as it is especially common on the road to the volcano of Kilauea. Dr. W. T. Brigham observed this rose more than forty years ago along the trail to the Kapena Falls on Oahu; and he believes it may have been taken to the other islands from this locality. It has not become naturalized on Oahu.

Rubus jamaicensis L.

Rubus jamaicensis L., Mant., i, 1767, 75.

Rubus jamaicensis Sw., In Griesb. Fl. Brit. W. Ind., 1864, 231.

The so-called thimble-berry is common on Hawaii and in certain places on Maui. It is said to have been introduced from the West Indies.

Dr. W. T. Brigham & C. N. Forbes, Kilauea, 1908. C. N. F. no. 238. M. Keanae, Maui.

LEGUMINOSEAE.

Caesalpinia sappan L.

Caesalpinia sappan L., Sp. Pl., 1753, 381.

Found in gulches near Wahiawa, Oahu.

C. N. F., no. 1443. O.

Cassia chamaecrista L.

Cassia chamaecrista L., Sp. Pl., 1753, 379.

Common about Honolulu, Oahu.

Heller, no. 1969.

Cassia laevigata Willd.

Cassia laevigata Willd., Enum. Hort. Berol., 1813, 441.

Occurs on all the islands. Rather common on Oahu.

Heller, no. 2295.

Crotalaria fulva Roxb.

Crotalaria fulva Roxb., Fl. Ind., 1832, 3, 266.

Crotalaria saltiana Andr.

Crotalaria saltiana Andr., Bot. Rep., 1811, pl. 648.

Crotalaria striata DC., Prod., 1825, ii, 131.

Crotalaria spectabilis Roth.

Crotalaria sericea Retz. (not Burm.), Obs. Bot., 1779-91, 3, 26.

Crotalaria spectabilis Roth., Nov. Pl. Sp., 1821, 341.

The Crotalarias form a conspicuous portion of the naturalized flora of the islands. [327]

Flemingia strobullifera R. Br.*Flemingia strobullifera* R. Br., Ait. Hort. Kew. Ed. ii, 1810-13, 350.*Flemingia bracteata* Wight, Ic., 1840-53, 268.*Flemingia fruticulosa* Wall, Cat., 1828, 5754.*Flemingia abrupta* Wall, Cat., 1828, 5755.

I am indebted to Mr. C. J. Austin for showing me this plant. It occurs in company with *Psidium guajava* at Nahiku, East Maui. The plant is widely spread in India, where it is said to be very variable. It is a small shrub, and can be distinguished by the conspicuous folded bracts which hide the small flowers.

Medicago apiculata Willd.*Medicago apiculata* Willd., Sp. Pl., iii, 1803, 1414.

Rather rare but probably has the same range as the next species. It can be recognized by the unarmed pods. In the Index Kewensis it is regarded as being synonymous with the next. C. N. F., no. 1718. O. Honolulu, Oahu.

Medicago denticulata Willd.*Medicago denticulata* Willd., Sp. Pl., iii, 1803, 1414.

The bur clover is more or less common on all the islands. According to Van Dine,⁵ it was introduced on Maui in 1882.

Medicago intertexta Mill.*Medicago intertexta* Mill., Gard. Dict. Ed. viii, 1768, n. 4.

The above species is reported to be common about Honolulu, by Heller.

Melilotus indica (L.) All.*Trifolium melilotus* L., Sp. Pl., 1753, 764.*Trifolium indicum* Thunb., Prod. Pl. Cap., 1800, 136.*Melilotus indica* All., Fl. Pedem I, 1785-89, 308.*Melilotus parviflora* Desf., Fl. Atlant., 1800, ii, 187.*Melilotus minima* Roth., Nov. Pl. Sp., 1821, 361.*Melilotus occidentalis* Nutt., ex Torr & Gray, Fl. N. Am., 1838, i, 321.*Melilotus regulosa* Willd., Enum. Hort. Berol., 1809, 789.

Rather common on Oahu and Hawaii; probably on all islands.

Trifolium repens L.*Trifolium repens* L., Sp. Pl., 1753, 767.

The white clover is common in the pastures on Haleakala, Maui.

GERANIACEAE.

Erodium cicutarium (L.) L'Her.*Geranium cicutarium* L., Sp. Pl., 1753, 680.*Erodium cicutarium* L'Her., Ait. Hort. Kew., 1789, 2, 414.

Oahu, not common.

⁵ Hawaii Agr. Exp. Station Rept., 1908, p. 25.

Geranium carolinianum L.*Geranium carolinianum* L., Sp. Pl., 1753, 683.

This plant not only occurs on Hawaii as stated by Hillebrand; but is also common on Haleakala, Maui.

C. N. F., no. 218. M.

MALVACEAE.

Modiola caroliniana (L.) Don.*Malva caroliniana* L., Sp. Pl., 1753, 688.*Modiola multifida* Moench. Meth., 1791, 620.*Modiola caroliniana* Don., Gen. Hist. Pl., 1831, i, 466.

Rather common on the slopes of Mauna Kea, Hawaii.

C. N. F., no. 458. H.

Two species of Abutilon not yet identified are very common on all the islands.

PASSIFLORIACEAE.

Passiflora edulis Sims.*Passiflora edulis* Sims., Bot. Mag., 1818, 1989.

Sometimes plentiful, and to be expected in the forests of all the islands. Hillebrand says in his flora, 1871, "already escaped into the woods of East Maui." Native name, *lilikoi*.

Passiflora foetida L.*Passiflora foetida* L., Sp. Pl., 1753, 959.

An occasional roadside plant.

ONAGRACEAE.

Epilobium Billardierianum ? Ser.*Epilobium Billardierianum* Ser., DC. Prod., 1843, iii, 41.

A species of Epilobium which I have provisionally referred to the above, is very common on Haleakala.

C. N. F., no. 173. M.

UMBELLIFERAE.

Apium leptophyllum (DC.) F. Mull.*Sison Ammi* L., Sp. Pl., 1753, 252.*Heliosciadium leptophyllum* DC., Prod., 1830, iv, 105.*Apium leptophyllum* F. Mull., Benth. Fl. Aust., 1866, iii, 372.*Apium Ammi* Urban., Mart. Fl. Bras., 1879, ii, pt. i, 341.

First observed at Makawao, Maui. Not common on the islands.
C. N. F., Makawao, Maui.

Aplum petroselinum* L.Apium petroselinum* L., Sp. Pl., 1753, 264.*Petroselinum sativum* Hoffm., Gen. Umb., 1814, 177.

Common parsley. A coarse leaved form occurs along irrigating ditches on West Maui. Reported from Hanapepe Valley, Kauai, by Heller.

C. N. F., no. 74. M. Wainee Gulch, Maui.

Foeniculum vulgare* (L.) Gaertn.Anethrum foeniculum* L., Sp. Pl., 1753, 263.*Foeniculum vulgare* Gaertn., Fruct. & Seem., 1788, i, 105.*Foeniculum foeniculum* Karst., Deutsch. Fl., 1880-83, 837.

The fennel is very common on Maui and in certain localities on Hawaii. Mentioned in Mann's Flora,⁶ but overlooked by Hillebrand.

PRIMULACEAE.***Anagallis arvensis* L.***Anagallis arvensis* L., Sp. Pl., 1753, 148.

Occurs on all the islands.

LOGANIACEAE.***Buddleia asiatica* Lour.***Buddleia asiatica* Lour., Fl. Cochinc., 72. Benth in DC. Prod., x, 416.

This plant is very common in the valleys of the Waianae range on Oahu, sometimes occurring as small trees six to ten feet high, while at other places it occurs as a subherbaceous plant of three to four feet. It also occurs in the dense forests of the Koolauloa range, and has all the appearances of an indigenous plant. One form is nearly glabrous.

C. N. F., Sept., 1908, Punaluu Mts. C. N. F., Makaha Valley, Kaala Mts., Feb. 12-19, 1909.

GENTIANACEAE.***Erythraea centaurium* (L.) Pers.***Gentiana centaurium* L., Sp. Pl., 1753, 229.*Erythraea centaurium* Pers., Syn., 1805, i, 283.

This plant is well established on East Maui and Hawaii. It is generally found on grazing land.

C. N. F., no. 180. M. Paia, Maui.

⁶Horace Mann. Flora of the Hawaiian Islands. Communications Essex Institute, Salem, Mass., vi, 1871.

CONVOLULACEAE.

Ipomoea chrysoides Ker-Gaul.

Ipomoea chrysoides Ker-Gaul., Bot. Reg., 270.

Nuuanu Valley, Oahu. Not common.

C. N. F., no. 1360. O.

Ipomoea grandiflora Lam.

Ipomoea glaberrima Boj., Hook. Journ. Bot., 1834, 357.

Ipomoea grandiflora Lam., Tab. Ency., i, 467. Roem. & Schult., Syst., iv, 240.

Dr. Guppy⁷ records his species for the first time from Hawaii. His localities were a mile west of Kapa-ahu and the coasts of Kalae. There are no specimens in this herbarium.

Ipomoea peltata Chois.

Convolvulus peltatus L., Sp. Pl., 1753, 121.

Ipomoea peltata Chois., Conv. Or., (?), 70. DC. Prod., ix, 359.

This species were observed by Mr. J. M. Lydgate a number of years ago in Wainiha Valley on Kauai, and judging from the locality in which he observed it, he had no hesitation in calling it indigenous. The flowers of our plant are yellow and all the leaves are peltate.

J. M. Lydgate, Wainiha Valley, Kauai, 1909.

LABIATAE.

Prunella vulgaris L.

Prunella vulgaris L., Sp. Pl., 1753, 600.

Very common on the cattle ranges of Haleakala, Maui. Also occurs on Hawaii but has not spread to any extent on this island.

SOLANACEAE.

Cestrum diurnum L.

Cestrum diurnum L., Sp. Pl., 1753, 191.

This species has escaped from cultivation and has become established in several places on Oahu. Several species of Cestrum have been in cultivation on these islands for a long time. I have heard the name Chinese ink-berry applied to this plant.

Nicotiana tabacum L.

Nicotiana tabacum L., Sp. Pl., 1753, 180.

Naturalized in various parts of Maui, especially on the north-east side.

⁷H. B. Guppy. Observations of a Naturalist in the Pacific, vol. ii, pp. 52, 554.

Solanum sodomeum* L.Solanum sodomeum* L., Sp. Pl., 1753, 187.

This African species is rather common on Oahu. First recorded by Heller.

ACANTHACEAE.***Thunbergia alata* Boj.***Thunbergia alata* Boj., Sims. Bot. Mag., 2591.

A troublesome garden weed in Honolulu.

***Thunbergia alata* var *aurantiaca* Ktz.**

This variety occurs as a roadside plant in Pauoa Valley on Oahu.

PLANTAGINACEAE.***Plantago lanceolata* L.***Plantago lanceolata* L., Sp. Pl., 1753, 113.

This species usually occurs at rather high elevations, 2000 to 3000 feet. Observed on all the islands.

RUBIACEAE.***Sherardia arvensis* L.***Sherardia arvensis* L., Sp. Pl., 1753, 102.

Maui, slopes of Haleakala.

C. N. F., no. 217. M.

CUCURBITACEAE.***Momordica charantia* L.***Momordica charantia* L., Sp. Pl., 1753, 1009.

Met with on all the islands, especially in rather dry localities where it climbs over rocks and stone walls. Near Kailua on Hawaii, it covers acres of rough aa flows, and is regarded as a bad pest.

Another cucurbit with spiny fruits not yet identified is fairly common on all the islands.

COMPOSITAE.***Carduus lanceolatus* L.***Carduus lanceolatus* L., Sp. Pl., 1753, 821.*Cirsium lanceolatum* Scop., Fl. Carn. Ed. ii, 1772, 2, 130.*Cnicus lanceolatus* Willd., Prodr. Fl. Berol., 1789, 259.

The common bur-thistle is well established on Hawaii and in several places on Maui.

C. N. F., no. 116. M. Hanaua W. Maui.

Chrysanthemum leucanthemum* L.Chrysanthemum leucanthemum* L., Sp. Pl., 1753, 888.*Leucanthemum vulgare* Lam., Fl. Fr., 1778, 2, 137.

The white daisy is rather common on Hawaii where it was observed growing on the slopes of Hualalai to the summit, and on the slopes of Mauna Loa.

Emilia flammea* Cass.Emilia flammea* Cass., Dict. Sc. Nat., 1819, 14, 406.

A roadside plant on Oahu, first recorded by Heller. There are specimens in the Museum herbarium collected several years before Heller's collection.

***Eupatorium* sp.**

A species of *Eupatorium* has lately become one of the most common plants on Maui.

C. N. F., no. 46. M. Hanakao, W. Maui. Dr. W. T. Brigham, Haleakala, E. Maui.

Hypochoeris radicata* L.Hypochoeris radicata* L., Sp. Pl., 1753, 811.

Rather common on Haleakala, Maui. Probably on Hawaii. Reported from Kauai by Heller.

Lapsana communis* L.Lapsana communis* L., Sp. Pl., 1753, 811.

Rather common on Maui and Hawaii.

Taraxacum officinale* (L.) Weber.Leontodon taraxacum* L., Sp. Pl., 1753, 798.*Taraxacum officinale* Weber, Prim. Pl. Holst., 1780, 56*Taraxacum dens-leonis* Desf., Fl. Atlant., 1800, 2, 228.*Taraxacum laraxacum* Karst., Deutsch. Fl., 1880-83, 1138.

The dandelion occurs on all the islands. It appears to have been introduced on the different islands from different sources. It has been observed on Kauai for at least twenty-five years, while a gentleman on Maui says it was introduced to that island very recently from New Zealand.

Zinnia pauciflora* L.Zinnia pauciflora* L., Sp. Pl., 1753, Ed. ii, 1269.

This species in company with *Verbesina enceloides* (Cav.) A. Gray, is very common about the cane fields in the vicinity of Lahaina, Maui.

C. N. F., Aug., 1910.

ADDENDA.**FICOIDEAE.**

Tetragonia expansa Murr.

Tetragonia expansa Murr., Comm. Gotting., 1783, vi, 13.

Tetragonia halimifolia Forst., Prod. 1786, 39.

Tetragonia cornuta Gaertn., Fruct., 1788, ii, 483.

Tetragonia inermis F. Mull., Linnaea., 1852, xxv, 384.

Tetragonia quadricornis Stokes, Bot. Mat. Med., (?), iii, 127.

This plant, generally known as the New Zealand Spinach was collected on the beach near Makapu point on Oahu. This is one of the localities where logs from the north-west coast of America are washed on to the beach. The plant is rather widely distributed on the shores of the Pacific, and has been cultivated in Europe.

C. N. F., no. 1406. O.

COMPOSITAE.

Senecio vulgaris L.

Senecio vulgaris L., Sp. Pl., 1753, 867.

The groundsel probably occurs on all the islands. It generally grows in partly forested sections. There are specimens in the herbarium from Maui, Hawaii and Oahu.

C. N. F., nos. 202. M. and 1695. O.

[334]

INDEX TO VOLUME IV.

The pagination of the volume will be found at the bottom of the page.
An asterisk denotes that the species or article is figured.

- Abudefduf amabilis*, 54.
 behnii, 53.
 caesio*, 53.
 coelestinus, 52.
 Jordani*, 53.
 leucopomus, 54.
 multifasciatus, 53.
 saxatilis, 54.
 septemfasciatus, 53.
 sordidus, 53.
 taupo, 54.
 uniocellatus, 54.
 zonatus, 55.
Abutilon sp. (2), 329.
Acanthochromis polyacanthus, 55.
Acanthurus garretti, 67.
 unicornis, 67.
Acridotheres tristis, 147.
Acrostichum calomelanos, 324.
Æstrelata sandwichensis, 137.
Alauda arvensis, 147.
Albula vulpes, 5.
Alticus saiensis, 88.
Amakihi, 160.
Amaranthus spinosus, 324.
Amblypistus tænianotus, 83.
Amia amboinensis, 32.
 arouensis, 33.
 novemfasciata, 32.
 savayensis, 33.
Anagallis arvensis, 330.
Anampses cæruleopunctatus, 55.
Anas wyvilliana, 144.
Anethrum sceniculum, 330.
Anguilla megastoma, 5.
Anomalops palpebratus, 20.
Anona cherimolia, 326.
Antennarius commersonii, 89.
 hispidus, 89.
*Antennarius leutescens**, 89.
Apapane, 158.
Apium ammi, 329.
 leptophyllum, 329.
 petroselinum, 330.
Arenaria rubra, 325.
Asio accipitrinus sandwichensis, 146.
Atherina lacunosa, 15.
Attendance of visitors, 1907, 100.
 1908, 182.
 1909, 233.
 1910, 305.
Auku killed by mongoose, 145.
Aulostomus valentini, 17.
Australian coll., W. R. Castle, 235.
Awaous crassilabris, 84.
 puntangoides, 84.
Balistes bursa, 73.
 capistratus, 74.
 chrysopterus, 73.
 flavimarginatus, 73.
 fuscus, 73.
 niger, 73.
 vidua, 73.
Balistapus aculeatus, 74.
 rectangulus, 74.
 undulatus, 74.
 verrucosus, 74.
Belone annulata, 12.
 platura, 12.
Blackman, L. G., resigns, 93.
Bone house at Moanalua, 130.
Borabora petroglyphs, 294.
Botanical accessions, 1909, 231.
 1910, 305.
Brassica campestris, 326.
Brighan, W. T.:—
 Annual Report, 1907, 93.

- Brigham, W. T.:—
 1908, 179.
 1909, 227.
 1910, 303.
 Curved adzes, 255.
- Bryan, W. A.:—
 almost drowned, 150.
 Birds of Molokai, 133.
 departs, 93.
 hunts the uau, 137.
- Buda marina, 325.
- Buddleia asiatica, 330.
- Cæsalpinia sappan, 327.
- Cæsio cœrulaureus, 44.
 lunaris, 44.
- Cæsio teres*, 45.
- Callyodon baliensis, 60.
 bataviensis, 60.
 celebicus, 60.
 cyanognathus, 60.
 dimidiatus, 60.
 dubius, 59.
 dussumieri, 62.
 erythrodon, 62.
 fasciatus, 62.
 lacerta, 60.
 lepidus, 60.
 macrocheilus, 60.
 moensi, 63.
 oviceps, 63.
 pentazonus, 60.
 pyrrhostethus, 60.
 quoyi, 62.
 strongylocephalus, 60.
 ultramarinus*, 63.
 unicolor, 60.
 waitei*, 60.
- Canthigaster compressus, 79.
 solandri, 79.
- Caranx forsteri, 32.
 lugubris, 32.
 melampygus, 32.
- Carcharias sorrah, 4.
- Carduus lanceolatus, 332.
- Carpodacus mexicanus, 148.
- Cartwright, B. Jr., loans collection, 309.
- Carved stone lamp, 127.
- Cassia chamaecrista, 327.
 Isevigate, 327.
- Casts of fishes, 113.
- Cephalopholis argus, 34.
 miniatus, 35.
 urodelus, 34.
- Ceremony at taking of fish, 210.
- Ceropterus calomelanos, 324.
 chrysophylla, 324.
- Cestrum diurnum, 331.
- Chætodon auriga, 64.
 citrinellus, 65.
 ephippium, 65.
 flavirostris, 64.
 gahhm, 66.
 lineolatus, 64.
 lunula, 64.
 ornatissimus, 65.
 pelewensis, 64.
 reticulatus, 65.
 semeion, 65.
 setifer, 64.
 ulietensis, 64.
 unimaculatus, 65.
 vagabundus, 64.
- Cheilinus digrammus, 58.
 fasciatus, 58.
 trilobatus, 59.
- Cheilio inermis, 57.
- Chlorodrepanis kalaana, 160.
- Chœrops macrodon, 55.
- Chromis axillaris, 55.
 cœruleus, 55.
- Chrysanthemum leucanthemum, 333.
- Cirsium lanceolatum, 332.
- Cnicus lanceolatus, 332.
- Conchological accessions, 1907, 104.
 1908, 186.
 1909, 231.
 1910, 304.
- Connelly collection, 228, 235.
- Convolvulus peltatus, 331.
- Cooke, C. M.: Accessions of Pulmonata, 231.

- Coris annulatus*, 57.
 ayula, 56.
 caudimaculatus, 57.
Coryphopterus criniger, 84.
Corythroichthys sealei, 17.
 waitei, 17.
Crepis japonica, 323.
Crotalaria saltiana, 327.
 sericea, 327.
 spectabilis, 327.
 striata, 327.
Ctenochaetus striatus, 67.
Curved adzes, Brigham, 255.
Cypselurus oligolepis, 15.
 *tahitensis**, 13.

Dascyllus aruanus, 55.
 melanurus, 55.
*Deleastes dæctor**, 81.
Deverill collection purchased, 99.
 list of, 183.
Director's report, 1907, 93.
 1908, 179.
 1909, 227.
 1810, 303.
Dole, S. B., leaves the Board, 179.
Drepanorhamphus funereus, 148.
Drymaria cordata, 325.

Echidna amblyodon, 12.
 polyzona, 12.
Eleotris fusca, 84.
Emilia flammula, 333.
Epibulus insidiator, 55.
Epilobium billardierianum, 329.
Epinephelus cæruleopunctatus, 36.
 corallicola, 35.
 dæmelii, 36.
 fasciatus, 36.
 maculatus, 35.
 merra, 35.
 pachycentrus, 36.
 undulosus, 35.
 *zapyrus**, 36.
Erodium cicutarium, 328.
Erysimum officinale, 326.

Erythraea centaurium, 330.
Ethnological accessions, 1907, 101.
 1908, 183.
 1909, 235.
 1910, 306.
Eupatorium sp., 333.
*Euphorbia rockii**, 214.
Europterygius xanthopterus, 11.
Exchange lists, 1907, 110.
 1908, 196.
 1909, 251.
 1910, 320.
*Exocarpus luteolus**, 296.
Exocætus volitans, 15.

Fish ponds, 199, 207.
Fish traps, Stokes, 199.
Fish weirs, 199.
 possible origin, 206.
Fishing by weirs, 199, 206.
Fishing, superstitions, 209, 211.
Fistularia depressa, 17.
 petimba, 17.
Flemingia abrupta, 328.
 bracteata, 328.
 fruticulosa, 328.
 strobulifera, 328.
Fræniculum vulgare, 330.
 fœniculum, 330.
Foot-prints cut in stone, 278, 286, 289.
Forbes, Chas. N.:—
 Appointed, 179.
 New plants, I, 213.
 New plants, II, 296.
 Naturalized plants, 323.
 report herbarium, 305.
Forcipiger longirostris, 64.
Fornander, Stokes index to, 180.
Pregata aquila, 144.
Fulica alai, 145.

Gallinula sandvicensis, 145.
Gentiana centaurium, 330.
Geological specimens, 237.
Geranium carolinianum, 329.
 cicutarium, 328.

- Gnathodentex aurolineatus*, 45.
Gobius oligolepis, 84.
 ornatus, 84.
Gomphosus tricolor, 58.
Grammistes sexfasciatus, 38.
Gymnogramma calomelanos, 324.
Gymnogramme calomelanos, 324.
Gymnothorax chalazius, 10.
 tactus, 6.
Gymnothorax favagineus, 7.
 flavomarginatus, 9.
 marquesensis*, 10.
 nebulosus, 6.
 nubilis, 7.
 pictus, 6.
 rhodocephalus, 6.
 tahitensis*, 7.
 tenebrosus, 7.
 thyroideus, 9.
 zonipectis*, 7.

Halichoeres centiquadrus, 56.
 modestus, 56.
 notophthalmus, 56.
 scapularis, 56.
 solorensis, 56.
 trimaculatus, 56.

Halophylla ovalis, 324.
 ovata, 324.

Hawaiian petroglyphs, 257, 284.
Heliosciadium leptophyllum, 329.
Hemigymnus melapterus, 55.

Hemiramphus affinis, 13.
 australensis*, 12.
 commersoni, 13.
 echanteri, 13.
 melanurus, 13.
 platurus, 13.

Heniochus chrysostomus, 65.
 permutatus, 65.

Hepatus achilles, 66.
 elongatus, 66.
 guttatus, 67.
 lineatus, 66.
 nigricans, 66.
 olivaceus, 66.
 triostegus, 66.

*Hesperomannia lydgatei**, 220.
Higgins, Miss E. B., appointed Librarian, 304.
Hilo petroglyphs, 278.
Himantopus knudseni, 145.
Himatione sanguinea, 158.
Hina, 203, 210.
Holacanthus cyanotis, 65.
 diacanthus, 66.

Holacanthus flavissimus, 65.
 imperator, 65.
 lineolatus, 65.
 loriculus, 65.
 permutatus, 65.

Holocentrus diadema, 24.
 diploxyphius, 24.
 ensifer, 20.
 lacteoguttatus, 20.
 microstomus, 24.
 opercularis, 24.
 pectilopterus, 24.
 praslin, 20.
 punctatissimus, 24.
 rubellio, 22.
 spinifer, 20.
 tiere, 24.
 tiereoides, 20.
 verticalis, 22.

Hologymnosus semidiscus, 57.
Holosteum cordatum, 325.

Hypochæris radicata, 333.

 Insect collections, 298.
Ipomæa cryseides, 331.
 glaberrima, 331.
 grandiflora, 331.
 peltata, 331.

Julis pulcherima, 57.

 Kakawahia, 164.
Kalaina wawae, 286.
Kamalalawalu, 259, 269, 273.
Kau petroglyphs, 273.
Kauai petroglyphs, 257, 291.
Kanalua, Kauwalua, 130.
Kohala petroglyphs, 279.

- Kona petroglyphs, 259.
 Kuhlia malo, 34.
 tæniura, 34.
 Kuula, 203, 209, 211.
 Kyphosus cinerascens, 47.

 Lapsana communis, 333.
 Lemna minor, 324.
 Leontodon taraxicum, 333.
 Lepidaplois axillaris, 55.
 Leptecheneis flaviventris*, 83.
 Lethrinus harak, 47.
 leutjanus, 47.
 mahsena, 47.
 moensi, 47.
 Leucanthemum vulgare, 333.
 Library accessions, 1907, 105.
 1908, 186.
 1909, 238.
 1910, 310.
 Liliuokalani loans, relics of chiefs, 309.
 Liza waigiensis, 15.
 Lo vulpinus*, 71.
 Loko, 199.
 Lophortyx californica, 146.
 Lutianus bengalensis, 43.
 bohar, 44.
 fulviflamma, 43.
 fulvus, 40.
 fuscescens, 43.
 gibbus, 41.
 marginatus, 41.
 melanesiae*, 43.
 monostigma, 40.
 nukuhivae*, 40.
 russellii, 43.
 semincinctus, 43.
 tahitiensis*, 40.
 Lysimachia longisepala*, 222.

 Macropharyngodon meleagris, 56.
 Malva caroliniana, 329.
 Mapo soporator, 84.
 Medicago apiculata 328.
 denticulata, 328.
 intertexta, 328.

 Megalops cyprinoides, 4.
 Melilotus indica, 328.
 minima, 328.
 parviflora, 328.
 regulosa, 328.
 Merinthe haplodactyla, 79.
 Micranous hawaiiensis, 137.
 Modiola caroliniana, 329.
 multifida, 329.

 Moho bishopi, 170.
 Molokai birds, Bryan, 133.
 Molokai petroglyphs, 257, 284, 290.
 Momordica charantia, 332.
 Monacanthus fatensis*, 74.
 spilosomus, 75.
 Monotaxis grandoculis, 46.
 Moorings drilled in boulders, 273.
 Moringua javanica, 12.
 Mugil cephalus, 17.
 kelaartii, 15.
 Mulloidess auriflamma, 48.
 flammeus, 48.
 ruber, 48.
 samoensis, 47.
 Munia nisoria, 148.
 Myripristis adustus, 26.
 microphthalmus, 28.
 murdjan, 26.
 phæopus*, 28.
 sanguineus*, 26.
 Myxus leuciscus, 15.

 New Hawaiian plants, Forbes, 213.
 Nicotiana tabacum, 331.
 Niihau petroglyphs, 293.
 Nycticorax nycticorax nævius, 145.

 Oahu petroglyphs, 257, 258, 290.
 Odontogobius phalæna, 84.
 Olomao, 171.
 Oreomystis flammea, 164, 167*.
 Ornithological accessions, 1907, 103.
 Ostracion auricauda*, 77.
 chryseres*, 75.
 cornutum, 75.
 dexter*, 75.

- Ostracion lentiginosum, 79.
 renardi, 77.
 sebae, 77.
 tuberculatum, 77.
- Ou, 166.
- Pakule, 201.
- Palmeria dolei, 158.
- Pa makiawa, 201.
- Paracanthistius maculatus, 35.
- Paracirrhites forsteri, 51.
- Paramia quinquelineata, 33.
- Parapercis xanthozena, 86.
- Pardachirus pavonius, 86.
- Passiflora edulis, 329.
 fetida, 329.
- Pelor didactylura, 83.
- Pempheris mangula, 32.
- Pentapus vittatus, 46.
- Periophthalmus barbarus, 84.
- Petroglyphs, Borabora, 294.
 cutting, 261.
 Hawaii, 2575, 284.
 Hilo, 278.
 Kau, 273.
 Kauai, 257, 291.
 Kohala, 279.
 Kona, Hawaii, 259.
 Molokai, 257, 284, 290.
 Niihau, 293.
 Oahu, 257, 290.
 Puna, 278.
 Stokes, J. F. G., 257.
 Tubuai, 293.
- Petroscirtes rhinorhynchus, 89.
 tapeinosoma, 89.
- Phaëornis rufa*, 171.
- Phaethon leptopus, 144.
- Phallic emblem, 266.
- Phallic stone ♀, 101.
- Pharopteryx nigricans, 38.
- Phasianus versicolor, 146.
 torquatus, 146.
- Plantago lanceolata, 332.
- Platax orbicularis, 63.
- Platophryns mancus, 86.
- Platyglossus marginatus, 56.
- Plotosus anguillaris, 12.
- Polycarpon tetraphyllum, 325.
- Polydactylus plebeius, 18.
- Pomacentrus cyanospilus, 52.
 lividus, 52.
 nigricans, 52.
 notopthalmus, 52.
 pavo, 52.
 scolopss, 52.
 teinurus, 52.
- Pomacentrus tripunctatus, 52.
- Portulaca flava, 325.
 lutea, 325.
- Priacanthus cruentatus, 38.
 hamruhr, 38.
- Pristiapogon, koilomatodon, 33.
 snyderi, 33.
- Promethichthys pacificus*, 28.
- Prunella vulgaris, 331.
- Pseudupeneus, aurantiacus, 48.
 barberinus, 49.
 bifasciatus, 51.
 chryserydros, 51.
 moana, 48.
 pleurospilos, 51.
 porphyreus, 51.
- Psittirostra psittacea, 166.
- Pterois antennata, 83.
- Pueo nesting, 146.
- Puffinus newelli, 143.
- Puna petroglyphs, 278.
- Querimana crenilabris, 15.
- Ranunculus hawaiiensis, 326.
 maniensis, 326.
 parviflorus, 326.
- Raphanus sativus, 326.
- Relics of chiefs, 238, 309.
- Reports, Director's, 1907, 93.
 1908, 179.
 1909, 227.
 1910, 303.
- Rosa sp., 327.
- Rubus jamaicensis, 327.
- Rumex acetosella, 324.
 patentia, 325.

- Salarias azureus*, 87.
cæsius, 88.
caudolineatus, 87.
coronarius, 86.
edentulus, 86.
hasselti, 84.
lineatus, 86.
marmoratus, 87.
meleagris, 86.
Salarias quadricornis, 86.
tubuensis, 87.
Sardinella kunzei, 5.
Saurida gracilis, 5.
Scarichthys auritus, 59.
coeruleopunctatus, 59.
rarotongæ, 59.
Schupp, Miss E.—
 appointed, 93, 98.
 resigns May, 1910, 304.
Scolopsis, *bilineata*, 45.
 monogramma, 45.
 temporalis, 45.
 trilineata, 45.
Scorpæna cooki, 79.
Scorpænopsis quiescens, 80.
Sculpture in relief, Stokes, 221.
Scuticaria marmorata, 12.
 tigrina, 11.
Seale, A., Fishes of S. Pacific, 3.
Sebastapistis, *baillieui*, 80.
 guamensis, 80.
 strongensis, 80.
Senecio vulgaris, 334.
Shark pen, 201, 212.
Sherardia arvensis, 332.
Siganus doliatetus, 69.
 marmoratus, 71.
 rostratus, 67.
 shortlandensis, 69.
 striolatus, 69.
 verrucosus, 69.
 zoniceps, 69.
Sison ammi, 329.
Sisymbrium officinale, 326.
 Skylark common on Molokai,
 147.
Solanum sodomeum, 332.
- Spergularia media*, 325.
salina, 325.
Spheroides hypselogenion, 79.
oblongus, 79.
Sphyræna goodingi, 18.
obtusata, 18.
Spilogelia chinesis, 146.
Stethojulis casturi, 56.
Stokes, J. F. G.:—
 Hawaiian Petroglyphs, 257.
 Stone sculpture in relief, 121.
 Walled Fish-traps of Pearl Harbor, 199.
Stolephorus delicatulus, 5.
Stone sculpture in relief, 121.
Sturtevant, H. F. dies, 93.
Superstitions of fishermen, 209, 211.
Swazey, O. H.:—
 Insect collections in Museum, 208.
 appointed, 93.
Synanceia verrucosa, 81.
Synodus varius, 5.
Taraxicum dens-leonis, 333.
officinale, 333.
Terapon jarbua, 45.
Tetragonia cornuta, 334.
 expansa, 334.
 halimifolia, 334.
 inermis, 334.
 quadricornis, 334.
Tetraodon aerostaticus, 79.
 immaculatus, 79.
 lachrymatus, 79.
 setosus, 79.
Thalassoma aneitense, 57.
 dorsale, 57.
 duperreyi, 57.
 fuscum, 57.
 lunare, 57.
 punctatum, 57.
 schwanefeldii, 57.
 umbrostigma, 57.
Thallurus chlorourus, 58.
Thunbergia alata, 332.
 aurantiaca, 332.
Tissa marina, 325.

- Trachinocephalus myops*, 5.
Trifolium indicum, 328.
 melilotus, 328.
 repens, 328.
Toxotes jaculator, 63.
Trachurops crumenophthalmus, 30.
Trachyrhamphus sp., 17.
Tubuai petroglyphs, 293.
Tylosurus choram, 12.
 giganteus, 12.

Uau, 137.
Upeneus vittatus, 51.
Uropterygius concolor, 11.

Variola leuti, 35.
Vestiaria coccinea, 155.
- Viola helena**, 218.
 *oahuensis**, 216.
Walled Fish-traps, 199.
Westervelt, Rev. W. D., gives Mexican antiquities, 304.

Xanthichthys rivulatus, 73.
Xystæma argyreum, 51.
*Yap money**, 102.

Zanclus canescens, 66.
Zebrasoma flavescens, 67.
 veliferum, 67.
Zinnia pauciflora, 333.

[342]