

New Cricket Genera and Species (Orthoptera: Grylloidea) from the Pacific Region Deposited in the Bishop Museum, Honolulu

DANIEL OTTE

Department of Entomology, Academy of Natural Sciences, Philadelphia, Pennsylvania 19103, USA; email: otte@ansp.org

Abstract: Ten new genera and 99 new species of crickets from Fiji and the western Pacific are validated through bibliographic reference to their original description and the addition of type depositories.

In two recent papers on Pacific crickets (Otte & Cowper, 2007; Otte, 2007) the type depositories were inadvertently omitted for the new species described, which resulted in them being invalid nomenclaturally and ultimately caused the new genus-group names based on them to also be invalid nomenclaturally. This paper provides a complete list of the new species giving bibliographic reference to characters to differentiate each species, indicates where the types are deposited, and validates 10 genus-group names with bibliographic references to characters differentiating each as well as giving a valid type species designation [NB: the genus *Nausorixipha* Otte & Cowper was validly proposed in Otte & Cowper (2007) as it was based on a previously described species].

The new species described from Fiji are currently deposited in the Bishop Museum in Honolulu (BPBM) but will eventually be deposited in the Fiji National Insect Collection in Suva (FNIC). The new species in Otte (2007) deriving from Bishop Museum material are permanently deposited in BPBM.

FIJI TAXA

Subfamily Gryllinae

Modicogryllus vaturu Otte & Cowper, n. sp.

Modicogryllus vaturu Otte & Cowper, 2007: 219.

Type.— Holotype male. FJ-92. FIJI: **Viti Levu**: Nausori Highlands, 8–14 km SE of jct of Vaturu Dam Rd and Sigatoka River Rd, 3.iii.1985, hand netted (Alexander, Otte and Rice) (FNIC; currently in BPBM).

Modicogryllus nandi Otte & Cowper, n. sp.

Modicogryllus nandi Otte & Cowper, 2007: 219.

Type.— Holotype male. FJ-88. FIJI: **Vanua Levu**: Labasa-Savusavu road, near mountain crest, rainforest, 23.ii.1985, hand netted (Alexander, Otte and Rice) (FNIC; currently in BPBM).

1. Contribution No. 2006-013 to the NSF-Fiji Arthropod Survey.

Modicogryllus volivoli Otte & Cowper, n. sp.*Modicogryllus volivoli* Otte & Cowper, 2007: 219.

Type.— Holotype male. FJ-81. FIJI: **Taveuni**: Cakaudrove, 3.2 km NW Lavena Vlg., Mt. Koronibuabua, 16.855°S, 179.892°W, 235 m, 5–17.vi.2004, Malaise 1 (E. I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Subfamily Nemobiinae

Dictyonemobius labasa Otte & Cowper, n. sp.*Dictyonemobius labasa* Otte & Cowper, 2007: 219.

Type.— Holotype male. FJ-93. FIJI: **Vanua Levu**: Labasa-Savusavu road, near mountain crest, rainforest, 23.ii.1985, hand netted (Alexander, Otte and Rice) (FNIC; currently in BPBM).

Dictyonemobius conaros Otte & Cowper, n. sp.*Dictyonemobius conaros* Otte & Cowper, 2007: 220.

Type.— Holotype male. FJ-84. FIJI: **Taveuni**: Cakaudrove, 5.3 km SE Tavuki Vlg., Mt. Devo, 16.841°S, 179.968°W, 1064m, 3–10.i.2003, Malaise 3 (E. I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Dictyonemobius savu Otte & Cowper, n. sp.*Dictyonemobius savu* Otte & Cowper, 2007: 220.

Type.— Holotype male. FJ-87. FIJI: **Viti Levu**: Vuda, 1 km E Abaca Vlg., Koroyanitu Ntl. Pk., Savuione Trail, 17°40'S, 177°33'E, 800 m, 7–12.x.2002, Malaise (E. I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Dictyonemobius trico Otte & Cowper, n. sp.*Dictyonemobius trico* Otte & Cowper, 2007: 220.

Type.— Holotype male. FJ-85. FIJI: **Taveuni**: Cakaudrove, 5.3 km SE Tavuki Vlg., Mt. Devo, 16.841°S, 179.968°W, 1064m, 3–10.i.2003, Malaise 3 (E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Subfamily Trigonidiinae

Trigonidium inopinum Otte & Cowper, n. sp.*Trigonidium inopinum* Otte & Cowper, 2007: 221.

Type.— Holotype male. FJ-61. FIJI: **Vanua Levu**: Labasa-Savusavu road, near mountain crest, rainforest, 23.ii.1985, hand netted (Alexander, Otte and Rice).

FIJIXIPHA Otte & Cowper, n. gen.*Fijixiphapha* Otte & Cowper, 2007: 221.

Type species.—*Fijixiphapha batia* Otte & Cowper, 2007, by present designation.

Fijixiphapha atalos Otte & Cowper, n. sp.*Fijixiphapha atalos* Otte & Cowper, 2007: 221.

Type.— Holotype male. FJ-33. FIJI: **Taveuni**: Cakaudrove, 5.3 km SE Tavuki Vlg., Mt. Devo, 16.841°S, 179.968°W, 1064 m, 2–10.x.2002, Malaise 3 (E. I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Fijixiphapha batia Otte & Cowper, n. sp.*Fijixiphapha batia* Otte & Cowper, 2007: 222.

Type.— Holotype male. FJ-26. FIJI: **Kadavu**: 0.25 km SW Solodamu Vlg., Moanakaka Bird Snctry, 19.078°S, 178.121°E, 60 m, 25.viii–23.x.2003, Malaise 4 (E. I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Fijixiphapha configens Otte & Cowper, n. sp.*Fijixiphapha configens* Otte & Cowper, 2007: 224.

Type.— Holotype male. FJ-56. FIJI: **Viti Levu**: Vuda, 1 km E Abaca Vlg., Koroyanitu Ntl. Pk., Kokabula Trail, 17°40'S, 177°33'E, 800 m, 12–19.xi.2002, Malaise (E. I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Fijixiphapha exuros Otte & Cowper, n. sp.*Fijixiphapha exuros* Otte & Cowper, 2007: 225.

Type.— Holotype male. FJ-41. FIJI: **Vanua Levu**: Bua, 6 km NW Kilaka, Batiqere Range, 16.8153°S, 178.9864°W, 146 m, 15.vi–28.vi.2004, Malaise (E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Fijixiphapha harpeza Otte & Cowper, n. sp.*Fijixiphapha harpeza* Otte & Cowper, 2007: 225.

Type.— Holotype male. FJ-1. FIJI: **Viti Levu**: Naitasiri, Nakobalevu Mt., rainforest, 18°03'S, 178°25'E, 340 m, 24.x–29.x.2003, Malaise (M.E. Irwin, E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Fijixiphapha inaudax Otte & Cowper, n. sp.*Fijixiphapha inaudax* Otte & Cowper, 2007: 225.

Type.— Holotype male. FJ-13. FIJI: **Kadavu**: 0.25 km SW Solodamu Vlg., Moanakaka Bird Snctry, 19°04'39"S, 178°07'15.6"E, 60 m, 19.xii.03–18.i.2004, Malaise (E.I. Schlinger, M. Tokota'a) [FBA 087423] (FNIC; currently in BPBM).

Fijixiphapha naitasiri Otte & Cowper, n. sp.*Fijixiphapha naitasiri* Otte & Cowper, 2007: 225.

Type.— Holotype male. FJ-8. FIJI: **Viti Levu**: Naitasiri, Navai Village, Eteni, 17°37'S, 177°59'E, 700 m, 15.v–2.vi.2003, Malaise (M.E. Irwin, E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Fijixiphapha penita Otte & Cowper, n. sp.*Fijixiphapha penita* Otte & Cowper, 2007: 226.

Type.— Holotype male. FJ-17. FIJI: **Vanua Levu**: Bua, Kilaka, 16°48'412"S, 178°59'290"E, 98 m, 28.vi–21.vii.2004, Malaise (M.E. Irwin, E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Fijixiphapha titilans Otte & Cowper, n. sp.*Fijixiphapha titilans* Otte & Cowper, 2007: 226.

Type.— Holotype male. FJ-42. FIJI: **Kadavu**: 0.25 km SW Solodamu Vlg., Moanakaka Bird Snctry, 19.078°S, 178.121°E, 60 m, 25.viii–23.x.2003, Malaise 3 (E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Fijixiphia valens Otte & Cowper, n. sp.*Fijixiphia valens* Otte & Cowper, 2007: 226.

Type.— Holotype male. FJ-45. FIJI: **Vanua Levu**: Macuata, Rokosalase, in forest, 16°31'891"S, 179°01'147"E, 105 m, 26.iii-9.iv.2004, Malaise (M.E. Irwin, E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Fijixiphia xipheres Otte & Cowper, n. sp.*Fijixiphia xipheres* Otte & Cowper, 2007: 226.

Type.— Holotype male. FJ-6. FIJI: **Viti Levu**: Naitasiri, 4 km WSW Colo-i-Suva Village, Mt. Nakobalevu, 18.055"S, 178.424"E, 372m, 14–28.vii.2003, Malaise 3 (E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

KADAVUXIPHA Otte & Cowper, n. gen.*Kadavuxiphia* Otte & Cowper, 2007: 227.

Type species.—*Kadavuxiphia soladamu* Otte & Cowper, 2007, by present designation.

Kadavuxiphia soladamu Otte & Cowper, n. sp.*Kadavuxiphia soladamu* Otte & Cowper, 2007: 227.

Type.— Holotype male. FIJI: **Kadavu**: Solodamu, coastal limestone forest, 19°04'S, 178°07'E, 128 m, 11.vi–6.vii.2003, Malaise (M.E. Irwin, E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

LEVUXIPHA Otte & Cowper, n. gen.*Levuxiphia* Otte & Cowper, 2007: 227.

Type species.—*Levuxiphia viticola* Otte & Cowper, 2007, by present designation.

Levuxiphia viticola Otte & Cowper, n. sp.*Levuxiphia viticola* Otte & Cowper, 2007: 227.

Type.— Holotype male. FJ-25. FIJI: **Viti Levu**: Naitasiri, 4 km WSW Colo-i-Suva Village, Mt. Nakobalevu, 18.055"S, 178.424"E, 372 m, 25.ii–17.iii.2003, Malaise 3 (E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

MINUTIXIPHA Otte & Cowper, n. gen.*Minutixiphia* Otte & Cowper, 2007: 228.

Type species.—*Minutixiphia exenios* Otte & Cowper, 2007, by present designation.

Minutixiphia exenios Otte & Cowper, n. sp.*Minutixiphia exenios* Otte & Cowper, 2007: 228.

Type.— Holotype male. FJ-62. FIJI: **Vanua Levu**: 12 km N of Savusavu, along lumber road, rainforest, 24.ii.1985, hand netted (Alexander, Otte and Rice) (FNIC; currently in BPBM).

Nausorixiphia dogotuki Otte & Cowper, n. sp.*Nausorixiphia dogotuki* Otte & Cowper, 2007: 228.

Type.— Holotype male. FJ-40. FIJI: **Vanua Levu**: Macuata, Dogotuki, 2.5 km E of Nasavu River, 16.2519"S, 179.7833"E, 226 m, 7.vii.2003, Malaise (M.E. Irwin, E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Nausorixipha euthetos* Otte & Cowper, n. sp.Nausorixipha euthetos* Otte & Cowper, 2007: 228.

Type.— Holotype male. FJ-29. FIJI: **Viti Levu:** Naitasiri, 3.2 km E Navai Vlg., Veilaselase Track, 17.624°S, 178.009°E, 1020 m, 6.vi–15.vii.2003, Malaise 2 (E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Nausorixipha nakubalevu* Otte & Cowper, n. sp.Nausorixipha nakubalevu* Otte & Cowper, 2007: 229.

Type.— Holotype male. FJ-22. FIJI: **Viti Levu:** Naitasiri, 4 km WSW Colo-i-Suva Village, Mt. Nakobalevu, 18.055°S, 178.424°E, 372 m, 25.ii–17.iii.2003, Malaise 3 (E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Nausorixipha navai* Otte & Cowper, n. sp.Nausorixipha navai* Otte & Cowper, 2007: 229.

Type.— Holotype male. FJ-10. FIJI: **Viti Levu:** Naitasiri, Navai Village, Eteni, 17°37'S, 177°59'E, 700 m, 15.v–2.vi.2003, Malaise (M.E. Irwin, E. I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Nausorixipha viti* Otte & Cowper, n. sp.Nausorixipha viti* Otte & Cowper, 2007: 229.

Type.— Holotype male. FJ-2. FIJI: **Viti Levu:** Sigatoka, Sigatoka Sand Dunes N.P., 18°09'939"S, 177°28'967"E, 4 m, 24.xi–16.xii.2003, Malaise (M.E. Irwin, E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

SAVUXIPHA* Otte & Cowper, n. gen.Savuxiphia* Otte & Cowper, 2007: 230.

Type species.—*Savuxiphia tromodes* Otte & Cowper, 2007, by present designation.

Savuxiphia tromodes* Otte & Cowper, n. sp.Savuxiphia tromodes* Otte & Cowper, 2007: 230.

Type.— Holotype male. FJ-52. FIJI: **Viti Levu:** Vuda, 1 km E Abaca Vlg., Koroyanitu Ntl. Pk, Savuione Trail, 17°40'S, 177°33'E, 800 m, 7–12.x.2002, Malaise (E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

TAVUKIXIPHA* Otte & Cowper, n. gen.Tavukixiphia* Otte & Cowper, 2007: 230.

Type species.—*Tavukixiphia devo* Otte & Cowper, 2007, by present designation.

Tavukixiphia devo* Otte & Cowper, n. sp.Tavukixiphia devo* Otte & Cowper, 2007: 230.

Type.— Holotype male. FIJI: **Vanua Levu:** Bua, Kilaka, 16°48'41.2"S, 178°59'29.0"E, 98 m, 28.vi–21.vii.2004, Malaise (M.E. Irwin, E. I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

VANUAXIPHA* Otte & Cowper, n. gen.Vanuaxiphia* Otte & Cowper, 2007: 230.

Type species.—*Vanuaxiphia perfixa* Otte & Cowper, 2007, by present designation.

Vanuaxiphad adamatos Otte & Cowper, n. sp.*Vanuaxiphad adamatos* Otte & Cowper, 2007: 230.

Type.— Holotype male. FJ-63. FIJI: **Viti Levu**: Nausori Highlands, 8–14 km SE of jct of Vaturu Dam Rd and Sigatoka River Rd, 3.iii.1985, hand netted (Alexander, Otte and Rice) (FNIC; currently in BPBM).

Vanuaxiphad dendriacos Otte & Cowper, n. sp.*Vanuaxiphad dendriacos* Otte & Cowper, 2007: 230.

Type.— Holotype male. FJ-54. FIJI: **Viti Levu**: Namosi, Ocean Pacific, Wainaddoi, coastal forest, 18°10'S, 178°15'E, 40 m, 5.xi–24.xi.2003, Malaise (M.E. Irwin, E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Vanuaxiphad eteni Otte & Cowper, n. sp.*Vanuaxiphad eteni* Otte & Cowper, 2007: 231.

Type.— Holotype male. FJ-49. FIJI: **Viti Levu**: Naitasiri, Navai Village, Eteni, 17°37'S, 177°59'E, 700 m, 13.ii–18.ii.2004, Malaise (M.E. Irwin, E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Vanuaxiphad koroni Otte & Cowper, n. sp.*Vanuaxiphad koroni* Otte & Cowper, 2007: 231.

Type.— Holotype male. FJ-37. FIJI: **Taveuni**: Cakaudrove, 3.2 km NW Lavena Vlg., Mt. Koronibuaabua, 16.855°S, 179.892°W, 235 m, 5–17.vi.2004, Malaise 1 (E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Vanuaxiphad perfixa Otte & Cowper, n. sp.*Vanuaxiphad perfixa* Otte & Cowper, 2007: 231.

Type.— Holotype male. FJ-58. FIJI: **Viti Levu**: Naitasiri, Navai Village, Eteni, 17°37'S, 177°59'E, 700 m, 24.x–8.xi.2003, Malaise (M.E. Irwin, E. I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Vanuaxiphad tricosa Otte & Cowper, n. sp.*Vanuaxiphad tricosa* Otte & Cowper, 2007: 231.

Type.— Holotype male. FJ-50. FIJI: **Viti Levu**: Namosi, Ocean Pacific, Wainaddoi, coastal forest, 18°10'S, 178°15'E, 40 m, 5.xi–24.xi.2003, Malaise (M.E. Irwin, E. I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Vanuaxiphad xylicos Otte & Cowper, n. sp.*Vanuaxiphad xylicos* Otte & Cowper, 2007: 232.

Type.— Holotype male. FIJI: **Vanua Levu**: Macuata, 0.6 km S of Rokosalase Village, forest, 16.5333°S, 179.0181°E, 180 m, 22.v–8.vi.2004, Malaise (E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

VEISARIXIPHA Otte & Cowper, n. gen.*Veisarixiphad* Otte & Cowper, 2007: 232.

Type species.—*Veisarixiphad waivudawa* Otte & Cowper, 2007, by present designation.

Veisarixiphia waivudawa* Otte & Cowper, n. sp.Veisarixiphia waivudawa* Otte & Cowper, 2007: 232.

Type.— Holotype male. FJ-4. FIJI: **Viti Levu**: Naitasiri, 3.8 km N Veisari Stlmt., log rd to Waivudawa, 18.079°S, 178.363°E, 300 m, 12.xii.2002–3.i.2003, Malaise 2 (E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

VITIXIPHA* Otte & Cowper, n. gen.Vitixiphia* Otte & Cowper, 2007: 232.

Type species.—*Vitixiphia axios* Otte & Cowper, 2007, by present designation.

Vitixiphia axios* Otte & Cowper, n. sp.Vitixiphia axios* Otte & Cowper, 2007: 232.

Type.— Holotype male. FJ-11. FIJI: **Taveuni**: Cakaudrove, 5.3 km SE Tavuki Vlg., Mt. Devo, 16.841°S, 179.968°W, 1064 m, 17–24.x.2002, Malaise 3 (E. I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Vitixiphia bua* Otte & Cowper, n. sp.Vitixiphia bua* Otte & Cowper, 2007: 233.

Type.— Holotype male. FJ-44. FIJI: **Vanua Levu**: Bua, 6 km NW Kilaka, Batiqere Range, 16.8153°S, 178.9864°W, 146 m, 15.vi–28.vi.2004, Malaise (E. I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Vitixiphia chlora* Otte & Cowper, n. sp.Vitixiphia chlora* Otte & Cowper, 2007: 233.

Type.— Holotype male. FJ-46. FIJI: **Taveuni**: Cakaudrove, Taveuni Estate, in garden, 16°50'S, 179°59'E, 140 m, 31.x–21.xi.2002, Malaise (M.E. Irwin, E. I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Vitixiphia kilaka* Otte & Cowper, n. sp.Vitixiphia kilaka* Otte & Cowper, 2007: 233.

Type.— Holotype male. FJ-15. FIJI: **Vanua Levu**: Bua, Kilaka, 16°48'412"S, 178°59'290"E, 98m, 28.vi–21.vii.2004, Malaise (M.E. Irwin, E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Vitixiphia peracta* Otte & Cowper, n. sp.Vitixiphia peracta* Otte & Cowper, 2007: 233.

Type.— Holotype male. FJ-59. FIJI: **Viti Levu**: Vuda, 1 km E Abaca Vlg., Koroyanitu Ntl. Pk, Savuione Trail, 17°40'S, 177°33'E, 800 m, 26.x–5.xi.2002, Malaise (E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Vitixiphia vuda* Otte & Cowper, n. sp.Vitixiphia vuda* Otte & Cowper, 2007: 233.

Type.— Holotype male. FJ-18. FIJI: **Viti Levu**: Vuda, Koroyanitu N.H.P., Abaca Village, 17°40'S, 177°33'E, 400 m, 6.v–26.v.2003, Malaise (M.E. Irwin, E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

VUDAXIPHA Otte & Cowper, n. gen.*Vudaxiphia* Otte & Cowper, 2007: 234.Type species.—*Vudaxiphia kokabula* Otte & Cowper, 2007, by present designation.*Vudaxiphia kokabula* Otte & Cowper, n. sp.*Vudaxiphia kokabula* Otte & Cowper, 2007: 234.Type.— Holotype male: FIJI: **Viti Levu**: Vuda, Koroyanitu Eco Park, 0.5 km N Abaca Vlg., 17.667°S, 177.55°E, 800 m, 12–19.xi.2002, Malaise 1 (E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Subfamily Podoscritinae

Aphonoides catastictos Otte & Cowper, n. sp.*Aphonoides catastictos* Otte & Cowper, 2007: 234.Type.— Holotype male. FJ-82. FIJI: **Viti Levu**: Naitasiri, 4 km WSW Colo-i-Suva Village, Mt. Nakobalevu, 18.055°S, 178.424°E, 372m, 14–28.vii.2003, Malaise 3 (E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).*Aphonoides kadavu* Otte & Cowper, n. sp.*Aphonoides kadavu* Otte & Cowper, 2007: 236.Type.— Holotype male. FJ-73. FIJI: **Kadavu**: Solodamu, coastal limestone forest, 19°04'S, 178°07'E, 128m, 23.x–19.xii.2003, Malaise (M.E. Irwin, E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).*Aphonoides namalata* Otte & Cowper , n. sp.*Aphonoides namalata* Otte & Cowper , 2007: 236.Type.— Holotype male. FJ-72. FIJI: **Viti Levu**: Vuda, Koroyanitu N.H.P. Abaca Village, 17°40'S, 177°33'E, 400 m, 6.v–26.v.2003, Malaise (M.E. Irwin, E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).*Aphonoides peristiges* Otte & Cowper, n. sp.*Aphonoides peristiges* Otte & Cowper, 2007: 237.Type.— Holotype male. FJ-76. FIJI: **Taveuni**: Cakaudrove, 5.3 km SE Tavuki, Devo Peak, 16.8431°S, 179.9681°W, 1064m, 14.xi–21.xi.2002, Malaise (E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).*Aphonoides tavuki* Otte & Cowper, n. sp.*Aphonoides tavuki* Otte & Cowper, 2007: 237.Type.— Holotype male. FJ-71. FIJI: **Taveuni**: Cakaudrove, 5.5 km SE of Tavuki Village, rainforest, 16.843°S, 179.966°W, 1188m, 30.vi–14.viii.2004, Malaise (E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).*Aphonoides xylurgos* Otte & Cowper, n. sp.*Aphonoides xylurgos* Otte & Cowper, 2007: 238.Type.— Holotype male. FJ-78. FIJI: **Vanua Levu**: Macuata, Rokosalase, in forest, 16°31'891"S, 179°01'147"E, 105 m, 26.iii–9.iv.2004, Malaise (M.E. Irwin, E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Subfamily Itarinae

Phalaria eugeris* Otte & Cowper, n. sp.Phalaria eugeris* Otte & Cowper, 2007: 238.

Type.— Holotype male. FJ-83. FIJI: Taveuni: Cakaudrove, 5.3 km SE Tavuki Village, Mt. Devo, 16°50'27.4"S, 179°58'04.1"W, 1064 m, 31.x–14.xi.2002, Malaise (E.I. Schlinger, M. Tokota'a) (FNIC; currently in BPBM).

Phalaria galoa* Otte & Cowper, n. sp.Phalaria galoa* Otte & Cowper, 2007: 238.

Type.— Holotype male. FJ-90. Viti Levu: Galoa Road ca. 10 N of Sigatoka-Suva Rd, rainforest, 25.ii.1985, hand netted (Alexander, Otte and Rice) (FNIC; currently in BPBM).

WESTERN PACIFIC TAXA

Cardiodactylus gressitti* Otte, n. sp.Cardiodactylus gressitti* Otte, 2007: 347.

Type.— Holotype male. SOLOMON ISLANDS: Mono Island, SE side, 50 m, 6–8 xi 1980 (Gressitt) (BPBM 16,760).

Cardiodactylus vella* Otte, n. sp.Cardiodactylus vella* Otte, 2007: 347.

Type.— Holotype male. SOLOMON ISLANDS: Vella Lavella, Mt. Arewana, 10–400 m, 16 xi 1963 (Gressitt) (BPBM 16,761).

Cardiodactylus sedlaceki* Otte, n. sp.Cardiodactylus sedlaceki* Otte, 2007: 347.

Type.— Holotype male. SOLOMON ISLANDS: San Cristobal, Wairahu River, 100–400 m, 9–15 v 1964 (J. Sedlacek) (BPBM 16,762).

Cardiodactylus maai* Otte, n. sp.Cardiodactylus maai* Otte, 2007: 347.

Type.— Holotype male. NEW GUINEA: Netherlands NG, River Tor (mouth) 4 km E of Hol Maffen, 2 vii 1959 (Maa) (BPBM 16,763).

Cardiodactylus quatei* Otte, n. sp.Cardiodactylus quatei* Otte, 2007: 347.

Type.— Holotype male. NEW GUINEA: West NG, Vogelkop, Kebar Valley, W of Manokwari, 550 m, 31 i 1962 (Quate and Quate) (BPBM 16,764).

Cardiodactylus kondoi* Otte, n. sp.Cardiodactylus kondoi* Otte, 2007: 348.

Type.— Holotype male. PHILIPPINES: (SW) Sulu: Tarawakan NE, 12 ii 1957 (Yoshio Kondo) (BPBM 16,765).

***Cardiodactylus riga* Otte, n. sp.**

Cardiodactylus riga Otte, 2007: 348.

Type.— Holotype male. PHILIPPINES: Camarines Sur, Mt. Iriga, 500–600 m, 3 iv 1962 (H.M. Torrevillas) (BPBM 16,766).

***Cardiodactylus brandti* Otte, n. sp.**

Cardiodactylus brandti Otte, 2007: 348.

Type.— Holotype male. NEW IRELAND: Schleinitz Mountains, Lelet Plateau, x 1959 (W.W. Brandt) (BPBM 16,767).

***Cardiodactylus manus* Otte, n. sp.**

Cardiodactylus manus Otte, 2007: 348.

Type.— Holotype male. MANUS ISLAND: Momote, 24 xii 1959 (Maa) (BPBM 16,768).

***Cardiodactylus pelagus* Otte, n. sp.**

Cardiodactylus pelagus Otte, 2007: 348.

Type.— Holotype male. BORNEO: Sarawak, Nanga Pelagus near Kapit, 180–585 m, 7–14 viii 1958 (Maa) (BPBM 16,769).

***Cardiodactylus muiri* Otte, n. sp.**

Cardiodactylus muiri Otte, 2007: 349.

Type.— Holotype male. CERAM: Piroe, ii 1909 (F. Muir) (BPBM 16,770).

***Cardiodactylus boharti* Otte, n. sp.**

Cardiodactylus boharti Otte, 2007: 349.

Type.— Holotype male. RYUKYU ISLANDS: Ishigaki Island, 10–15 xi 1952 (G.E. Bohart) (BPBM 16,771).

***Cardiodactylus kraussi* Otte, n. sp.**

Cardiodactylus kraussi Otte, 2007: 349.

Type.— Holotype male. TONGA: ‘Eua Island, Ohonua, ii 1956 (N.L.H. Krauss) (BPBM 16,772).

***Cardiodactylus esakii* Otte, n. sp.**

Cardiodactylus esakii Otte, 2007: 349.

Type.— Holotype male. PALAU: Angaur: Saipan-Kitamura, 26 ii 1938 (Teiso Esaki) (BPBM 16,773).

***Cardiodactylus murakami* Otte, n. sp.**

Cardiodactylus murakami Otte, 2007: 349.

Type.— Holotype male. PALAU: Peliliou: Garukyoku, 20 i 1938 (Shiro Murakami) (BPBM 16,774).

Cardiodactylus hentownesi* Otte, n. sp.Cardiodactylus hentownesi* Otte, 2007: 350.

Type.— Holotype male. PONAPE: Mt. Dolennankap, 1800 feet, 11 viii 1946 (H.K. Townes) (BPBM 16,775).

Cardiodactylus kusaiense* Otte, n. sp.Cardiodactylus kusaiense* Otte, 2007: 350.

Type.— Holotype male. KUSAIE ISLAND: Mt. Tafeyat, 1000–1200 feet, 20 viii 1946 (H.K. Townes) (BPBM 16,776).

Cardiodactylus gagnei* Otte, n. sp.Cardiodactylus gagnei* Otte, 2007: 350.Type.— Holotype male. VANUATU: Ambrym, Ranon to Mt. Toyo, 0–500 m, 2 ix 1979 (W.C. Gagné *et al.*) (BPBM 16,777).***Cardiodactylus cheesmani* Otte, n. sp.***Cardiodactylus cheesmani* Otte, 2007: 350.

Type.— Holotype male. VANUATU: Tanna, x 1930 (L.E. Cheesman) (BPBM 16,778).

Cardiodactylus bulolo* Otte, n. sp.Cardiodactylus bulolo* Otte, 2007: 350.

Type.— Holotype male. NEW GUINEA: (NE) Bulolo, 700 m, 26 xi 1969 (Sedlacek) (BPBM 16,779).

Cardiodactylus mamai* Otte, n. sp.Cardiodactylus mamai* Otte, 2007: 350.

Type.— Holotype male. NEW GUINEA: (SE), Mamai Plantation, E of Port Glasgow, 150 m, 31 i 1965 (BPBM 16,780).

Cardiodactylus empagatao* Otte, n. sp.Cardiodactylus empagatao* Otte, 2007: 351.

Type.— Holotype male. PHILIPPINES: Misamis Or., Mt. Empagatao, 28 km SW of Gingog, 12 iv 1961 (H.M. Torrevillas) (BPBM 16,781).

Cardiodactylus efordi* Otte, n. sp.Cardiodactylus efordi* Otte, 2007: 351.

Type.— Holotype male. SOLOMON ISLANDS: Bougainville (S.), Kokure, near Crown Prince Range, 900 m, 8 vi 1966 (E.J. Ford Jr) (BPBM 16,782).

Cardiodactylus busu* Otte, n. sp.Cardiodactylus busu* Otte, 2007: 351.

Type.— Holotype male. NEW GUINEA: (NW) Japen Island, SSE of Sumberbaba, Dawai River, 10 x 1962 (N. Wilson) (BPBM 16,783).

Cardiodactylus tangtalau* Otte, n. sp.Cardiodactylus tangtalau* Otte, 2007: 353.

Type.— Holotype male. SOLOMON ISLANDS: Malaita, Tangtalau, 200 m, 26 ix 1957 (BPBM 16,784).

***Cardiodactylus kokure* Otte, n. sp.**

Cardiodactylus kokure Otte, 2007: 353.

Type.— Holotype male. SOLOMON ISLANDS: Buka Agricultural Station, 6–10 xii 1959 (T.C. Maa) (BPBM 16,785).

***Cardiodactylus kuschei* Otte, n. sp.**

Cardiodactylus kuschei Otte, 2007: 353.

Type.— Holotype male. SOLOMON ISLANDS: Guadalcanal, 2-5-21 (Kusche) (BPBM 16,786).

***Cardiodactylus mumurai* Otte, n. sp.**

Cardiodactylus mumurai Otte, 2007: 353.

Type.— Holotype male. SOLOMON ISLANDS: Bougainville, Mumurai, 9 vi 1956 (Gressitt) (BPBM 16,787).

***Cardiodactylus shanahani* Otte, n. sp.**

Cardiodactylus shanahani Otte, 2007: 353.

Type.— Holotype male. SOLOMON ISLANDS: Kolombangara, Pepele, 30 m, 7 ii 1964 (Shanahan) (BPBM 16,788).

***Cardiodactylus enkraussi* Otte, n. sp.**

Cardiodactylus enkraussi Otte, 2007: 353.

Type.— Holotype male. VANUATU: Santo, Segond Channel, viii 1950 (Krauss) (BPBM 16,789).

***Cardiodactylus tathimani* Otte, n. sp.**

Cardiodactylus tathimani Otte, 2007: 354.

Type.— Holotype male. SOLOMON ISLANDS: NW Malaita Island, Dala, 11 vi 1964 (Straatman) (BPBM 16,790).

***Cardiodactylus wairahu* Otte, n. sp.**

Cardiodactylus wairahu Otte, 2007: 354.

Type.— Holotype male. SOLOMON ISLANDS: San Cristobal, Wairahu River, 100 m, 9–15 v 1964 (Sedlacek) (BPBM 16,791).

***Cardiodactylus kukugai* Otte, n. sp.**

Cardiodactylus kukugai Otte, 2007: 354.

Type.— Holotype male. SOLOMON ISLANDS: Bougainville, Kukugai Village, 150 m, x 1960 (W.W. Brandt) (BPBM 16,792).

***Cardiodactylus javarere* Otte, n. sp.**

Cardiodactylus javarere Otte, 2007: 354.

Type.— Holotype male. NEW GUINEA: Papua, Daradae, near Javarere, Musgrove River, 100 m?, 3 x 1958 (Gressitt) (BPBM 16,793).

***Cardiodactylus togerao* Otte, n. sp.**

Cardiodactylus togerao Otte, 2007: 354.

Type.— Holotype male. SOLOMON ISLANDS: Bougainville (E), Togerao, 600 m, 15 iv 1968 (Straatman) (BPBM 16,794).

***Cardiodactylus singuawa* Otte, n. sp.**

Cardiodactylus singuawa Otte, 2007: 355.

Type.— Holotype male. NEW GUINEA: (NE) Lae, Singuawa River, 147°10'E 6°45'S, 30 m, 1 iv 1966 (G. Lippert) (BPBM 16,795).

***Cardiodactylus kolombangara* Otte, n. sp.**

Cardiodactylus kolombangara Otte, 2007: 355.

Type.— Holotype male. SOLOMON ISLANDS: Kolombangara, Gollifer's Camp, 700 m, 24 i 1964 (Shanahan) (BPBM 16,796).

***Cardiodactylus lavella* Otte, n. sp.**

Cardiodactylus lavella Otte, 2007: 355.

Type.— Holotype male. SOLOMON ISLANDS: Vella Lavella, Ulo Crater, 10 m, xii 1963 (Shanahan) (BPBM 16,797).

***Madasumma zamboanga* Otte, n. sp.**

Madasumma zamboanga Otte, 2007: 355.

Type.— Holotype male. PHILIPPINE ISLANDS: Mindanao, Zamboanga del Norte, Masawan Trail to Mount Malindang, 1290 m, 14 vii 1958 (H. Milliron) (BPBM 16,798).

***Lebinthus makilingus* Otte, n. sp.**

Lebinthus makilingus Otte, 2007: 356.

Type.— Holotype male. PHILIPPINE ISLANDS: Luzon Island, Mount Makiling, 400 m, 16 v 1959 (BPBM 16,799).

***Landreva erromanga* Otte, n. sp.**

Landreva erromanga Otte, 2007: 356.

Type.— Holotype male. VANUATU: Erromanga Island, August 1930 (L.E. Cheeseman) (BPBM 16,800).

***Mikluchomaklaia chuaye* Otte, n. sp.**

Mikluchomaklaia chuaye Otte, 2007: 356.

Type.— Holotype male. NEW GUINEA: Chuaye, 2000 m, 8–14 iii 1968 (BPBM 16,801).

***Mikluchomaklaia biaru* Otte, n. sp.**

Mikluchomaklaia biaru Otte, 2007: 356.

Type.— Holotype male. NEW GUINEA (NE): Upper Biaru, 1600–2200 m, 10 iii 1971 (Gressitt and Tawi) (BPBM 16,802).

***Mikluchomaklaia gorochovi* Otte, n. sp.**

Mikluchomaklaia gorochovi Otte, 2007: 356.

Type.— Holotype male. NEW GUINEA (NE): Upper Biaru, 1600–2200 m, 10 iii 1971 (Gressitt and Tawi) (BPBM 16,803).

***Modicogryllus abrichtos* Otte, n. sp.**

Modicogryllus abrichtos Otte, 2007: 357.

Type.— Holotype male. FJ-95. SOLOMON ISLANDS: Guadalcanal Honiara region, Mt Austin area forest edge, 5.iii.1985, hand netted (Alexander, Otte & Rice) (BPBM 16,804).

LITERATURE CITED

- Otte, D. 2007. New species of *Cardiodactylus* from the western Pacific region. *Proceedings of the Academy of Natural Sciences of Philadelphia* **156**: 341–400.
—. & G. Cowper. 2007. New cricket species from the Fiji Islands (Orthoptera: Grylloidea). *Proceedings of the Academy of Natural Sciences of Philadelphia* **156**: 217–303.