

## 38. Family SCENOPINIDAE\*

By Lewis P. Kelsey

The family Scenopinidae is distributed worldwide and currently includes nearly 400 described species. The Australasian/Oceanian fauna contains 86 species, mostly from Australia. Six of the 17 known genera are found in the Australasian/Oceanian Regions, including 2 worldwide genera (*Scenopinus*, *Metatrachia*) and 4 genera endemic to Australia (*Neopseudotrichia*, *Paramonova*, *Paratrichia*, *Riekiella*).

Most of the species represented have a rather limited distributional pattern restricted at or near the type locality. However, there are 4 species with wide distributions undoubtedly associated with the movement of trade goods (*Scenopinus fenestralis* and *S. glabifrons* are cosmopolitan, *S. lucidus* is from Egypt, Nigeria, and the Hawaiian Islands, and *S. papuanus* is from Hawaii, Taiwan, New Guinea, and Sri Lanka).

Very little is known of the larval biology except that they seem to be predators of other insects. Larvae have been found associated with stored-grain pests, in nests of birds and rodents, in burrows of beetle larvae attacking a rather wide range of trees and shrubs, and in soil in association with therevid larvae, which they closely resemble. Adults are taken in the field by sweeping flowers with open nectaries that are accessible to the sponging mouthparts of the adult. In buildings they are taken at windows, giving rise to the common name "window flies" for species of this family.

Ref.: Kelsey (1969, 1970a,b, 1987, rev., keys), Paramonov (1955g, syst., biol.), Krivosheina (1980, biol.).

### Genus METATRICHIA Coquillett

**METATRICHIA** Coquillett, 1900d: 500. Type species: *Scenopinus bulbosa* Osten Sacken, 1877, orig. des.

*papuana* Kelsey, 1970a: 44. PNG (PNG).  
*waterhousei* Paramonov, 1955g: 640 (*Pseudomphrale*). Australia (NSW).

### Genus NEOPSEUDOTRICHIA Kelsey

**NEOPSEUDOTRICHIA** Kelsey, 1969: 299. Type species: *Neopseudotrichia kewi* Kelsey, 1969, orig. des. *canei* Kelsey, 1969: 200. Australia (SA).

*mariaensis*, Paramonov, 1955, not Hardy, 1933, misid.  
*inglewoodi* Kelsey, 1970b: 141. Australia (Qld).  
*kewi* Kelsey, 1969: 300. Australia (ACT); Australia (NSW).  
*mariaensis* Hardy, 1933b: 419 (*Pseudotrichia*). Australia (Tas). N. comb.  
*monteithi* Kelsey, 1970b: 144. Australia (Qld); Australia (NSW).

### Genus PARAMONOVA Kelsey

**PARAMONOVA** Kelsey, 1970b: 125. Type species: *Riekiella quadrifida* Kelsey, 1969, orig. des.  
*arila* Kelsey, 1987: 10. Australia (Qld).  
*brevipalpa* Kelsey, 1970b: 127. Australia (NSW).  
*bungabandii* Kelsey, 1975: 205. Australia (WA).  
*burrijunana* Kelsey, 1987: 11. Australia (WA).  
*bushi* Kelsey, 1971: 203. Australia (WA).  
*collessi* Kelsey, 1969: 307 (*Riekiella*). Australia (NSW). N. comb.

\*Published as Miscellaneous Paper No. 1146 of the Delaware Agricultural Experiment Station, Contribution No. 573 of the Department of Entomology and Applied Ecology, University of Delaware, Newark, Delaware.

**complexa** Kelsey, 1969: 309 (*Riekiella*). Australia (ACT). **N. comb.**  
**dycei** Kelsey, 1969: 310 (*Riekiella*). Australia (NSW). **N. comb.**  
**emarginata** Kelsey, 1970b: 129 (*Riekiella*). Australia (WA).  
**fonsamnicula** Kelsey, 1980: 9. Australia (NSW).  
**hamata** Kelsey, 1980: 10. Australia (NSW).  
**mangara** Kelsey, 1987: 12. Australia (WA).  
**poocherania** Kelsey, 1987: 13. Australia (SA).  
**quadrifida** Kelsey, 1969: 313 (*Riekiella*). Australia (Qld). **N. comb.**  
**taylori** Kelsey, 1969: 315 (*Riekiella*). Australia (Vic). **N. comb.**  
**uncata** Kelsey, 1969: 315 (*Riekiella*). Australia (NSW). **N. comb.**  
**zentae** Kelsey, 1975: 207. Australia (SA).

#### Genus PARATRICHIA Kelsey

**PARATRICHIA** Kelsey, 1969: 320. Type species: *Paratrichia lobosa* Kelsey, 1969, orig. des.  
**lobosa** Kelsey, 1969: 320. Australia (ACT).  
**spicata** Kelsey, 1975: 208. Australia (WA).  
**westralica** Paramonov, 1955g: 419 (*Pseudomphrale*). Australia (WA). **N. comb.**

#### Genus RIEKIELLA Paramonov

**RIEKIELLA** Paramonov, 1955g: 652. Type species: *Riekiella bicornis* Paramonov, 1955, orig. des.  
**arilaringa** Kelsey, 1987: 16. Australia (WA).  
**arnheimeri** Kelsey, 1975: 210. Australia (NT).  
**bicornis** Paramonov, 1955g: 653. Australia (Qld).  
**binnaliweanda** Kelsey, 1987: 18. Australia (WA).  
**bolgarti** Kelsey, 1969: 55 (*Scenopinus*). Australia (WA). **N. comb.**  
**burririmba** Kelsey, 1987: 19. Australia (NT).  
**cetratra** Kelsey, 1971: 196. Australia (NT).  
**cornuta** Kelsey, 1971: 197. Australia (NT).  
**culbara** Kelsey, 1987: 20. Australia (NT).  
**cuspidata** Kelsey, 1971: 198. Australia (Qld).  
**denticulata** Kelsey, 1971: 200. Australia (NT).  
**kilyalpa** Kelsey, 1987: 21. Australia (WA).  
**longitermina** Kelsey, 1975: 211. Australia (WA).  
**longiventris** Kelsey, 1969: 312. Australia (WA).  
**malkana** Kelsey, 1987: 22. Australia (WA).  
**maloga** Kelsey, 1987: 23. Australia (WA).  
**murilla** Kelsey, 1987: 24. Australia (WA).  
**nancyae** Kelsey, 1987: 25. Australia (WA).  
**nicholsi** Kelsey, 1971: 200. Australia (NT).  
**noona** Kelsey, 1987: 27. Australia (WA).  
**rossi** Kelsey, 1969: 123. Australia (WA).  
**rupari** Kelsey, 1987: 27. Australia (WA).  
**tamala** Kelsey, 1987: 29. Australia (WA).  
**woodwardi** Kelsey, 1971: 201. Australia (NT).  
**yoota** Kelsey, 1987: 30. Australia (WA).

#### Genus SCENOPINUS Latreille

**OMPIIRALE** Meigen, 1800: 29. Suppressed by I.C.Z.N., 1963b: 339.  
**SCENOPINUS** Latreille, 1802: 463. Type species: *Musca fenestralis* Linnaeus, 1758, mon.  
**SCENOPOEUS** Agassiz, 1847: 333 (unjust. emend. of *Scenopinus*).  
**LUCIDOMPHRALE** Kröber, 1937: 219. Type species: *Scenopinus lucida* Becker, 1902, orig. des.

- PAROMPHRALE* Kröber, 1937: 222. Type species: *Scenopinus glabrifrons* Meigen, 1824, orig. des.
- SCENOPINULA* Paramonov, 1955g: 651. Type species: *Scenopinula pallidipennis* Paramonov, 1955, mon.
- SCAENOPIUS*, error for *Scenopinus*.
- adventicius* Hardy, 1960: 329. Hawaiian Is.
- adyeri* Kelsey, 1987: 5. Australia (WA).
- arrectus* Kelsey, 1975: 198. Australia (WA)
- ascensus* Kelsey, 1987: 6. Australia (Qld).
- bicuspidis* Kelsey, 1971: 185. Australia (NT).
- biroi* Kertész, 1899b: 173. PNG (PNG); Australia (NSW, Qld).
- brittoni* Kelsey, 1970: 111. Australia (WA).
- casurarinus* Kelsey, 1975: 200. Australia (NT).
- cristafrons* Kelsey, 1971: 186. Australia (NT).
- darwini* Kelsey, 1969: 64. Australia (NT).
- dilkara* Kelsey, 1987: 7. Australia (WA).
- erugatus* Kelsey, 1975: 201. Australia (WA).
- fenestralis* Linnaeus, 1758: 597 (*Musca*). "Europa"; Australia (Qld); cosmopol.
- perkinsi*, Paramonov, 1955, not Hardy, 1942, misid.
- fijianus* Kröber, 1939a: 399 (*Omphrale*). Fiji.
- fragosifrons* Kelsey, 1969: 68. Australia (WA).
- glabrifrons* Meigen, 1824: 114. Austria; Australia (NSW, Qld, WA); cosmopol.
- civiculus* Hardy, 1933b: 418. Australia (NSW & Qld & SA).
- harleyi* Kelsey, 1971: 187. Australia (Qld).
- limbunayi* Kelsey, 1971: 190. Australia (NT).
- lucidus* Becker, 1902: 39. Egypt; Hawaiian Is; Algeria, Christmas I (Indian Ocean), Afrotrop. Reg.
- lunulanotafrons* Kelsey, 1971: 191. Australia (NT).
- mendumae* Kelsey, 1971: 188. Australia (NT).
- millarae* Kelsey, 1971: 189. Australia (Qld).
- minutus* Kelsey, 1969: 78. Australia (NSW).
- nakkari* Kelsey, 1987: 8. Australia (WA).
- norrisi* Kelsey, 1969: 82. Australia (Qld).
- pallidipennis* Paramonov, 1955g: 651 (*Scenopinula*). Australia (NSW); Australia (SA). N. comb.
- papuanus* Kröber, 1912e: 25 (*Omphrale*). PNG (PNG) & Taiwan; Guam, Hawaiian Is, Northern Marianas; Christmas I (Indian Ocean), Cocos (Keeling) Is, Sri Lanka.
- zeylanicus* Senior-White, 1922a: 205. Sri Lanka.
- niger*, Grimshaw, 1901, not De Geer, 1776, misid.
- perkinsi* Hardy, 1942: 202. Australia (Qld); Australia (NSW).
- renneri* Kelsey, 1975: 204. Australia (NT).
- sanfordi* Kelsey, 1971: 192. Australia (NT).
- spurrelli* Kelsey, 1969: 90. Australia (Vic).
- torrensianus* Kelsey, 1980: 7. Australia (SA).
- trigelasinus* Kelsey, 1971: 194. Australia (NT).
- triquetus* Kelsey, 1971: 192. Australia (NT).